



# Impact of a Head Covering on Photo-Therapy Induced Hypo-calcaemia in Full-term Neonates with Hyper-bilirubinemia

Sachin Mulye <sup>a\*</sup>, Revatdhama J. Meshram <sup>b<sup>o</sup></sup> and Krishnakumar Thakrani <sup>c<sup>#</sup></sup>

<sup>a</sup> Department of Neonatology, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha-442001, India.

<sup>b</sup> Department of Pediatrics, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Sawangi (Meghe), Wardha-442001, India.

<sup>c</sup> Department of Pediatrics, Datta Meghe Medical College, Nagpur-441110, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JPRI/2021/v33i58B34177

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/67577>

**Original Research Article**

**Received 07 October 2021**

**Accepted 14 December 2021**

**Published 15 December 2021**

## **ABSTRACT**

**Background:** Hyper-bilirubinemia causes yellowish discoloration of the sclera and skin in newborns and is one of the most common disorders that neonatologists face daily. In the first week of life, 60 percent of term and 80 percent of preterm babies have jaundice. In neonates with unconjugated hyper-bilirubinemia, phototherapy is the most common treatment.

**Objective:** To compare the occurrence of hypocalcemia in full-term icteric neonates with and without head covering during phototherapy.

**Materials and Methodology:** Two hundred and eight full-term neonates were split into two groups for photo-therapy for hyper-bilirubinemia. The heads of the participants in the study group were covered with a hat. Before beginning photo-therapy, serum bilirubin and serum calcium levels were assessed, as well as after 48 hours of photo-therapy. SPSS software was used to tabulate and analyze the data.

**Results:** Ninety-four (45%) of the 208 neonates in the sample were males, while 114 (55%) were

<sup>\*</sup> Assistant Professor

<sup>o</sup> Associate Professor

<sup>#</sup> Senior Resident

females. The average age was  $5.56 \pm 2.78$  days, and the average weight was  $3.04 \pm 0.58$  kilograms. Out of 208 neonates, 44 (21%) had hypocalcemia. Hypocalcemia occurred in 28 neonates in the control group and 16 neonates in the study group. There was no symptomatic hypocalcemia in any of the newborns. The p-value for the Chi square test was 0.2. Since this value is greater than 0.05, it is not statistically important.

**Conclusion:** In comparison to neonates whose heads were not covered, fewer neonates with covered heads experienced hypocalcemia in our research. However, these findings were not statistically important. However, Research in other countries found that neonatal head cover at phototherapy had a substantial impact. As a result, we propose that a major multicenter study be performed in our country to determine the exact significance.

*Keywords: Head covering; hyper-bilirubinemia; hypocalcemia; neonatal jaundice; photo-therapy.*

## 1. INTRODUCTION

It is normal for newborns to develop jaundice in their first few days of life. Jaundice affects 55 percent of full-term neonates around the world, according to reports. According to a study performed at a neonatal center, 28 percent of babies presented with jaundice [1]. This incidence was estimated at 37 percent in another study conducted in Swat. High bilirubin levels are usually harmless, but in a limited number of neonates, they can cause acute bilirubin encephalopathy, which can lead to kernicterus. Kernicterus affects 0.5 to 1.3 neonates per 1 lakh live births in developed countries, while it affects 3 percent of all neonates admitted to hospitals in developing countries [2,3]. Untreated hyperbilirubinemia can cause severe complications that can be avoided with early detection. These neonates are mostly treated with photo-therapy, with only a few requiring exchange transfusions [4,5]. The aim of treatment is to lower the level of unconjugated bilirubin to a safe level. With new and improved phototherapy equipment and techniques, the need for exchange transfusions has reduced.

Phototherapy is associated with few risks, including hyperthermia, rash, diarrhea, dehydration, DNA damage, retinal damage, bronze baby syndrome, patent ductus arteriosus in preterm infants, and calcium deficiency [6-8]. In babies, persistent hypo-calcemia can cause convulsions, apnea, and tetany. Cortisol causes hypocalcemia by increasing calcium absorption in the bones. Melatonin, a hormone produced by the pineal gland, blocks the effects of cortisol on bones [9]. The pineal secretion of melatonin is blocked during phototherapy, and the effect of cortisol is unopposed, resulting in hypocalcemia. Some researchers recommend intravenous calcium to prevent hypocalcemia during phototherapy, but rapid infusion in neonates may

cause cardiac arrest or hypotension [10,11]. As a result, other strategies for preventing hypocalcemia should be investigated [12]. Hypocalcemia can be avoided by shielding the heads of neonates during photo-therapy, according to a study conducted in Iran. They found that the sample group (covered heads) had a lower rate of hypocalcemia (14%) than the control group (uncovered heads) at the end of the study (39 %). To our knowledge, there hasn't been much research on this subject in India [13,14]. If we can avoid hypocalcemia in neonates by covering their heads during photo-therapy, we will not only use this technique in our neonatal unit, but we will also campaign for it in other hospitals to save babies from photo-therapy-induced hypocalcemia.

During phototherapy, we wanted to see how head covering affected ionized serum calcium and magnesium levels in neonates with hyperbilirubinemia.

### 1.1 Objective

To compare the occurrence of hypocalcemia in full-term icteric neonates with and without head covering during photo-therapy.

## 2. METHODOLOGY

This one-year prospective, comparative, case control study took place in the neonatal intensive care unit of a neonatology department at a medical college in Central India. In each category, the sample size was determined to be 104. There was a total of 208 neonates in the study. A non-probability (consecutive) sampling method was used. The study included neonates with a gestational age of more than 37 weeks, a weight of more than 2.5 kg, normal serum calcium levels before phototherapy, and hyperbilirubinemia (as defined by the operational

definition). Since any of these conditions can alter serum calcium levels, neonates with a history or evidence of asphyxia, sepsis, exchange transfusion, hemolytic anaemia, or hypothyroidism were excluded. Neonates were assessed using inclusion and exclusion criteria. The research was explained to the parents, and they signed a written informed consent form. Full-term stable neonates were randomly divided into two groups using computer-generated codes that were placed in envelopes. During photo-therapy, a stockinette cap was used to cover the heads of neonates in one group (study group), while the heads of neonates in the other group (control group) were not covered and were exposed. Sex, weight, serum total bilirubin level, and serum calcium level were documented before starting photo-therapy. Full blood count and reticulocyte count ruled out hemolytic anaemia, and C-reactive protein ruled out sepsis. Photo-therapy with four 40-watt blue lamps with a wave length of 420-470 nanometers was started on a regular basis. The neonate was held at a 40-centimeter distance from the lamps. The serum calcium level was measured 48 hours post initiation of photo-therapy. SPSS version 2.0 was used to evaluate the data collected. For qualitative variables like gender and the frequency of hypocalcemia, as well as quantitative variables like age, calcium, and bilirubin levels, frequencies and percentages were determined. Stratification was used to regulate effect modifiers such as age and gender. The chi-square test was used after stratification. A p-value of less than 0.05 was considered significant.

### 2.1 Operational Definitions

Neonate: Age range: 0 to 28 days. Hypocalcemia is a condition in which the body's calcium levels are low Total serum calcium 1.9

mmol/L in neonates aged 0-10 days, and total serum calcium 2.25 mmol/L in neonates aged >10 days, estimated 48 hours after photo-therapy commencement. Hyper-bilirubinemia was defined as serum total bilirubin levels of >188mol/L on day one, >257mol/L on day two, >306mol/L on day three, and >342mol/L on day four.

### 3. RESULTS

A total of 208 term neonates with hyper-bilirubinemia were included in this study. Male neonates made up 45 percent of the total, while female neonates made up 55 percent. Their average age was 5.56 ±2.78 days, and their average weight was 3.02±0.58 kilograms. The demographic data in both groups was identical, as shown in Table 1.

The mean serum total bilirubin level in all neonates was 381.32 ±45.31 mmol/L before photo-therapy, and 268.29 ± 46.46 mmol/L after 48 hours of photo-therapy. Before and after photo-therapy, the levels in the control group were 379.21 ± 43.4 mmol/L and 266.60 ±48.38 mmol/L, respectively. Before and after photo-therapy, the study group had 382.45 ±48.60 mmol/L and 270.43 ±45.48 mmol/L, respectively.

Similarly, before starting photo-therapy, all neonates' mean serum calcium was 2.40 ±0.28 mmol/L, and after 48 hours, it was 2.08 ±0.28 mmol/L. Before and after photo-therapy, the levels in the control group were 2.36 ±0.28 mmol/L and 1.48 ±0.24 mmol/L, respectively. In the study group, it was 2.46 ±0.24 mmol/L before photo-therapy and 2.18 ±0.26 mmol/L after photo-therapy, respectively. In both groups, serum calcium levels decreased, but the decrease in serum calcium levels was greater in the control group.

**Table 1. Demographic data of patients**

Gender	Control	Study
Male	44	50
Female	60	54
Age (Days)	4.38 ± 1.42	5.56 ± 2.78
Weight (Kg)	2.50 ± 0.70	3.04 ± 0.58

**Table 2. Frequency of hypocalcemia after photo-therapy**

Hypo-calcemia	Control	Study	p-value
Occurred	28 (27 %)	16 (15 %)	
Not occurred	76 (73 %)	88 (85 %)	0.2

**Table 3. Stratification of hypocalcemia with or without covering the head with respect to age**

Age Group	Hypo-calcemia	Control	Study	Total	p-value
<3 days	Yes	6	2	8	0.2
	No	20	26	46	
>3 days	Yes	22	14	36	0.3
	No	56	62	118	
Total		104	104	208	

**Table 4. Stratification of hypocalcemia with or without covering the head with respect to Gender**

Age Group	Hypo-calcemia	Control	Study	Total	p-value
Male	Yes	10	10	20	0.8
	No	34	40	74	
Female	Yes	18	6	24	0.08
	No	42	48	90	
Total		104	104	208	

Out of 208 neonates, 44 (21%) had hypocalcemia. There was no symptomatic hypocalcemia in any of the newborns. Hypocalcemia found in 28 neonates in the control group and 16 neonates in the study group. Table 2 provides more details. Tables 3 and 4 demonstrate the age and gender stratification of hypocalcemia with and without head covering.

#### 4. DISCUSSION

Phototherapy is a relatively safe and effective way to lower bilirubin levels in newborns with jaundice. However, complications such as hypocalcemia may occur. Hypocalcemia may also be accompanied by jitteriness, irritability, and lethargy. The neonates in the control group received photo-therapy in the traditional manner. Hypocalcemia affected 27% of neonates, but they were asymptomatic [15]. In term neonates, a study published in 2015 found similar findings. After photo-therapy, 30% of term neonates experienced hypo-calcemia, according to the report. A higher percentage of neonates experienced hypo-calcemia and were symptomatic in a few other trials. Hypocalcemia in neonates undergoing photo-therapy was studied in a research conducted in India. According to the findings, 67 % of neonates experienced hypocalcemia, with 81 percent of hypocalcemic neonates being symptomatic [16]. Similarly, after phototherapy, 66 % of term neonates experienced hypocalcemia, according to another study.

However, a small number of studies claim that photo-therapy has no effect on calcium levels in neonates with hyper-bilirubinemia. There have

been a few experiments in which the head was covered to minimize the sensitivity of the pineal gland to photo-therapy, resulting in lower hypocalcemia. When neonates' heads were covered, they experienced a substantial decrease in hypocalcemia. In Egypt, an analysis involving 124 neonates was carried out. They found that 24 % of neonates in the group without hats and 9.7% of neonates in the hat group experienced hypocalcemia [17]. After 48 hours of phototherapy, 24% of term neonates with exposed heads and 11% of term neonates with covered heads experienced hypocalcemia, according to another study conducted in Egypt. In a related study from Iran, 72 full-term newborns were included. According to their findings, 39% of neonates with an exposed head and 14% of neonates with a covered head experienced hypocalcemia. Hypocalcemia found in 27% of the neonates in our sample who had their heads exposed. This percentage is close to that found in Egyptian studies, but lower than that found in Iranian studies [18]. Fifteen percent (15%) of the neonates in our sample who had their heads covered developed hypocalcemia. This result is similar to that of an Iranian study, but it is higher than that of an Egyptian study. Despite the fact that our research showed a reduction in hypocalcemia by covering the heads of neonates, our findings were statistically insignificant in comparison to all of the previous studies that had shown substantial results. Another research in Iran found that covering the heads of icteric preterm infants during photo-therapy reduced hypocalcemia significantly. In this sample, the control group had hypocalcemia in 53% of the babies, while the head covered group in just 6% of the neonates.

Another Iranian research found that covering the heads of neonates during phototherapy reduced hypocalcemia significantly [19]. According to the t-test, the difference between the two classes was important ( $p=0.03$ ). During the analysis, we encountered a few limitations. To ensure that the neonates' heads were protected at all times during phototherapy, even during feeding, close contact with them was needed. Observing symptomatic hypocalcemia necessitates close supervision. However, due to a staffing crisis, this was not always possible. The research was carried out in a single centre with a limited sample size. To better understand the importance of head covering, further research should be performed in several centers with a broad sample size.

## 5. CONCLUSION

Many studies have shown that phototherapy causes hypocalcemia, and a few studies have shown a substantial reduction in hypocalcemia when the neonate's head is covered. While our analysis found that head covering reduced hypocalcemia, the results were statistically insignificant. To assess the exact importance of this intervention, more broad studies should be performed. If significant, it is a simple and inexpensive way to prevent hypo-calcemia in neonates undergoing photo-therapy.

## CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

The study was conducted after getting approval by the hospital's ethics review committee.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Brits H, Adendorff J, Huisamen D, Beukes D, Botha K, Herbst H, Joubert G. The prevalence of neonatal jaundice and risk factors in healthy term neonates at National District Hospital in Bloemfontein. *Afr J Prim Health Care Fam Med.* 2018; 10(1): a1582.
2. Tikmani SS, Warraich HJ, Abbasi F, Rizvi A, Darmstadt GL, Zaidi AKM. Incidence of neonatal hyperbilirubinemia: a population-based prospective study in Pakistan. *Trop Med Int Health.* 2010;15:502-07.
3. Haq UI, Israrulhaq, Khan S, Sayed Z. Common etiological spectrum of indirect hyperbilirubinemia in neonates. *JSMC.* 2017; 7(2):112-16.
4. Alkén J, Håkansson S, Ekéus C, Gustafson P, Norman M. Rates of extreme neonatal hyperbilirubinemia and kernicterus in children and adherence to national guidelines for screening, diagnosis, and treatment in Sweden. *JAMA Netw Open.* 2019; 2(3):e190858.
5. Nakade M, Jungari ML, Ambad R, Dhingra G. Status of Vitamins and Minerals in Pregnancy: Still A Point of Concern in Central India. *Int J Cur Res Rev.* July. 2020;12(14); 45-49.
6. Ebbesen F, Hansen TWR, Maisels MJ. Update on Phototherapy in Jaundiced Neonates. *Curr Pediatr Rev.* 2017; 13(3):176-80.
7. Abdelazeem KS, Soliman AA, Askar EAA. Efficacy of intensive phototherapy in management of neonatal hyperbilirubinemia in neonatal unit of Assiut university children Hospital. *J Neonatal Biol.* 2017; 6:266.
8. Singh PK, Chaudhuri PK, Chaudhuri AK. Phototherapy induced hypocalcemia in neonatal hyperbilirubinemia. *IOSR-JDMS.* 2017;16: 35-38.
9. Goyal S, Srivastava A, Bhattacharjee P, Goyal I, Malhotra K. Effect of phototherapy on serum calcium levels in neonates receiving phototherapy for neonatal jaundice. *Int J Res Med Sci.* 2018; 6(6):1992-95.
10. Kargar M, Jamshidi Z, Beheshtipour N, Pishva N, Jamali M. Effect of head covering on phototherapy-induced hypocalcaemia in icterus newborns; A randomized controlled trial. *Intl J Community Based Midwifery.* 2014; 2(2):121-26.
11. Ambad R, Dhok A. The association of lipid profile and uric acid levels in normotensive, preeclamptic pregnancy – A hospital based study. *J Datta Meghe Inst Med Sci Univ.* 2020;15:21-5.
12. Prabhakar N, Lazarus M, Ahirwar M. Effect of phototherapy on serum ionic calcium level in neonates with hyperbilirubinemia. *Ejpmr.* 2016; 3(4): 510-14.

13. Yadav RK, Sethi RS, Sethi AS. The evaluation of the effect of phototherapy on serum calcium level. *People's J Sci Res.* 2012; 5(2):1-4.
14. Srinivasa S, Renukananda S. Effect of phototherapy on hypo-calcemia. *J of Evolution Med Dent Sci.* 2015;4(24): 4165-68.
15. Paymaneh AT, Sajjadian N, Eivazzadeh B. Prevalence of photo-therapy induced hypocalcemia in term neonate. *Iran J Pediatr.* 2013;23(6):710-11.
16. Ezzeldin Z, Mansi Y, Abdelhamid TA, Sabry M. The effect of hat on phototherapy-induced hypocalcemia in jaundiced full-term neonates. *Dove Press.* 2015;5:73-78.
17. Nouh MA, El-saeed WF, Shehata AE, Mostafa SA. Impact of covering the heads of icteric neonates during phototherapy on the prevalence of hypocalcemia. *Med j Cairo Univ.* 2013;81(2):219-22.
18. Samane ZD. The effect of covering head on the hypocalcemia caused by phototherapy in the icteric preterm infants in the Vali-e-Asr Hospital in 2015: a randomized controlled trial. *Intl J Med Res Health Sci.* 2016;5:138-42.
19. Berekatain B, Badiea Z, Hoseini N. The effect of head covering in prevention of phototherapy-induced hypocalcemia in icterus new borns with gestational age less than 35 weeks. *Adv Biomed Res.* 2016; 5:176.

© 2021 Mulye et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/67577>