



Successful Ross Procedure for Aortic Valve Dysfunction in a Young Patient with Laubry-pezzi Syndrome: A Case with Ischemic Stroke as a Post-operative Complication

L. Afendi ^{a*}, L. Laklalech ^a, F. Essadqi ^a, A. El Bouazizi ^a,
M. Bouziane ^a and A, Drighil ^a

^a *Department of Cardiology, CHU Ibn Rochd, Casablanca, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/120431>

Case Report

Received: 26/05/2024

Accepted: 29/07/2024

Published: 03/08/2024

ABSTRACT

The article discusses a 21-year-old patient with Laubry-Pezzi syndrome who underwent a Ross procedure due to dysfunction of a mechanical aortic valve originally implanted for a congenital malformation. The Ross procedure, using the patient's own pulmonary valve, offers advantages for young patients. Despite successful valve replacement, the patient experienced a post-operative ischemic stroke which resolved with anticoagulation. The case underscores the importance of a multidisciplinary approach and close monitoring for optimal outcomes, contributing valuable insights to the management of similar conditions.

*Corresponding author: E-mail: afendilamiaa@gmail.com;

Cite as: Afendi, L., L. Laklalech, F. Essadqi, A. El Bouazizi, M. Bouziane, and A, Drighil. 2024. "Successful Ross Procedure for Aortic Valve Dysfunction in a Young Patient With Laubry-Pezzi Syndrome: A Case With Ischemic Stroke As a Post-Operative Complication". *Asian Journal of Research in Cardiovascular Diseases* 6 (1):51-55. <https://journalijrcc.com/index.php/AJRCD/article/view/102>.

Keywords: Laubrey-pezzi syndrome; congenital disease; cardiac abnormalities.

1. INTRODUCTION

Laubrey-Pezzi syndrome is a rare and complex congenital disease characterized by structural cardiac abnormalities, often including a ventricular septal defect (VSD) and valvular dysfunctions. Management of this syndrome requires a multidisciplinary approach and repeated surgical interventions throughout the patient's life due to the progressive nature of the cardiac anomalies [1].

The Ross procedure, which involves replacing the diseased aortic valve with the patient's own pulmonary valve and implanting a bioprosthesis in the pulmonary position, offers several advantages. It uses the patient's own tissues, reducing the risks of rejection and thrombosis, and improves hemodynamic outcomes [2]. This technique is particularly beneficial for young patients as it adapts to growth and increased metabolic needs [3].

Here, we present the case of a 21-year-old patient with Laubrey-Pezzi syndrome who underwent multiple surgical interventions, including a Ross procedure due to dysfunction of a mechanical aortic valve. This case illustrates the challenges and complications associated with managing this complex syndrome [4].

2. CASE PRESENTATION

The patient known with Laubrey-Pezzi syndrome since birth, underwent surgical closure for a ventricular septal defect (VSD) at the age of 1 years old. At 20 years old, he underwent

reoperation for an aortic valve replacement with implantation of a mechanical valve. In the immediate postoperative period, the patient presented with valve mismatch with high transaortic gradients of 65 mmHg, indicating dysfunction of the mechanical valve, necessitating a Ross procedure. At admission for the Ross procedure, the patient was asymptomatic except for exertional dyspnea. Vital signs were stable (BP 148/80 mmHg, HR 90 bpm) and ECG showed a regular sinus rhythm with negative T waves in the anteroseptal-apical territory. Echocardiography revealed a dilated non-hypertrophied left ventricle with mild dysfunction and an ejection fraction of 47% (Fig. 1), a non-dilated right ventricle with dysfunction, and dysfunction of the aortic valve prosthesis with a mean gradient of 65 mmHg (Fig. 2) and moderate paravalvular leak (Fig. 3). Pulmonary artery pressure was 32 mmHg. The work up showed a microcytic hypochromic anemia with a haemoglobin of 11g/dL and a CRP of 10 mg/L. After the Ross procedure, during which the diseased aortic valve was replaced with the patient's own pulmonary valve and a bioprosthesis was placed in the pulmonary position, the patient developed muscle weakness in the left upper limb. An emergency brain CT scan revealed a right frontal hypodensity of ischemic origin. Curative anticoagulation was initiated. The patient recovered muscle strength, and the follow-up CT.

scan was normal. Transthoracic echocardiography performed five weeks after surgery showed normal transaortic and transpulmonary gradients (Fig. 4).



Fig. 1. Apical 4-Chamber Echocardiographic View Showing A Dilated Dysfunctional Left Ventricle With FEVG: 47%




Fig. 2. Continuous doppler on the aortic prosthesis showing a high mean transprosthetic gradient of 65 mmHg




Fig. 3. Moderate aortic paraprosthetic leak




Fig. 4. Post-operative evaluation of the aortic valve: parasternal long-axis and short axis views

3. DISCUSSION

The Ross procedure, developed by Donald Ross in 1967, is a complex surgical technique but particularly beneficial for young patients requiring aortic valve replacement [5]. This technique involves harvesting the patient's pulmonary valve and implanting it in the aortic position, with a bioprosthesis placed in the pulmonary position. Using the patient's own tissues to replace the aortic valve offers several advantages, including better adaptation to hemodynamic needs, reduced risks of rejection and thrombosis, and increased durability [6]. The autologous pulmonary valve can better tolerate the high systemic pressures of the aortic circulation, thereby reducing the risks of long-term complications [2]. Prosthesis-patient mismatch is a well-documented complication of aortic valve replacements, where the size of the prosthesis is inadequate relative to the patient's aortic annulus, leading to high transvalvular gradients and valve dysfunction [7]. In the reported case, the patient presented with a transaortic gradient of 65 mmHg after the placement of a mechanical valve, indicating severe mismatch. This problem is critical as it can increase the risk of cardiac complications, including heart failure and prosthetic failure [8]. The Ross procedure, although more complex, offers an effective solution to this problem by using autologous tissues, thus reducing the risks of mismatch and thromboembolic complications [3].

The development of a cerebrovascular accident (CVA) after cardiac surgery is a dreaded complication, although rare, especially in a young patient of 21 years. The incidence of postoperative CVA varies, but factors such as aortic manipulations, emboli from thrombus or prosthetic material, and episodes of cerebral hypoperfusion can increase this risk [9]. In the presented case, the patient developed muscle weakness in the left upper limb, diagnosed as a right frontal hypodensity of ischemic origin on a brain CT scan. Management of this complication involved initiating curative anticoagulation, which allowed complete recovery of muscle strength and normalization of images on follow-up CT scan.

Prevention of neurological complications relies on close monitoring of high-risk patients, optimization of surgical and anesthetic techniques, and rapid management of postoperative neurological signs [10].

4. CONCLUSION

This case illustrates the clinical challenges and potential complications in managing patients with Laubrey-Pezzi syndrome requiring multiple valvular interventions. The Ross procedure appears to be a viable solution for young patients, despite the associated risks. Close postoperative monitoring and a multidisciplinary management approach are essential to optimize long-term outcomes. The rarity of this case and the complexity of the medical and surgical management underscore the value of this report. By documenting specific challenges and applied solutions, this article contributes to enriching the scientific literature on Laubrey-Pezzi syndrome and postoperative complications of the Ross procedure, providing valuable information for the future management of patients with similar conditions.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. El-Hamamsy I, Yacoub MH. The ross operation: A comprehensive review. *Circ.* 2009;119(6):840-849.
2. Pibarot P, Dumesnil JG. Prosthesis-patient mismatch in the aortic valve position. *Am J Cardiol.* 2000;86(6):459-460.
3. Yoganathan AP, et al. Aortic valve hemodynamics after the Ross procedure

- and homograft implantation: An *In vitro* study. *Ann Thorac Surg.* 2001;71(4):1061-1068.
Available:<https://pubmed.ncbi.nlm.nih.gov/11308182>.
4. Elkins RC, et al. The Ross operation: 16-year experience. *J Thorac Cardiovasc Surg.* 1999;117(5):667-680.
 5. Bove T, et al. Ross operation in young adults: Over 20 years of experience. *Eur J Cardiothorac Surg.* 2011;40(6):1480-1486.
 6. Kouchoukos NT, et al. The Ross procedure: Long-term clinical and echocardiographic follow-up. *Ann Thorac Surg.* 2004;78(3):773-781.
 7. Bacha EA, et al. The Ross procedure in neonates and infants: Is it a good solution for complex left ventricular outflow tract obstruction? *J Thorac Cardiovasc Surg.* 2001;122(2):261-270.
 8. Meyns B et al. The Ross operation: A word of caution. *Circulation.* 2000;102(19):III1-III4.
 9. da Costa FD, et al. Long-term results of the Ross operation: Analysis of 46 consecutive patients. *Ann Thorac Surg.* 2002;73(2):609-614.
 10. Roach GW, et al. Cerebral complications following cardiac surgery: The multicenter study of perioperative ischemia research group and the ischemia research and education foundation investigators. *Stroke.* 1996;27(6):1185-1191.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/120431>