



## **Prevalence and Distribution of Urogenital Schistosomiasis and Trichomoniasis in Oju LGA, Benue State, Nigeria**

**J. I. Chikwendu<sup>1\*</sup>, T. S. Atsuwe<sup>1</sup>, V. U. Obisike<sup>1</sup> and O. E. Igbor<sup>2</sup>**

<sup>1</sup>University of Agriculture, Makurdi, Benue State, Nigeria.

<sup>2</sup>Benue State University, Benue State, Nigeria.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors JIC and OEI designed the study, carried out the field research and wrote the manuscript. Authors TSA and VUO managed the literature search and performed the statistical analysis. All authors read and approved the final manuscript.*

### **Article Information**

#### Editor(s):

(1) Chiang-Ming Chen, Associate Professor, Department of Economics, University Rd., Puli, Nantou, 54561, R.O.C. National Chi Nan University, Taiwan.

#### Reviewers:

(1) Joseph Mukuni, School of Education, Virginia Tech, USA.

(2) Iroju Olaronke, Adeyemi College of Education, Ondo, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46858>

**Original Research Article**

**Received 25 April 2019**

**Accepted 06 July 2019**

**Published 15 July 2019**

### **ABSTRACT**

Schistosomiasis is a disease of medical importance. Schistosomiasis is prevalent in Nigeria where it is responsible for grave economic losses. A cross-sectional study was conducted from July, 2017-December, 2018 among school children from six (6) schools in Oju local government area of Benue State Nigeria, in order to determine the prevalence of urinary schistosomiasis with respect to sex, age and location. A total of 300 samples were collected and examined with the aid of a compound microscope (X10) and (X40) for the presence of *Schistosoma haematobium* eggs. The mean age of children that participated in the study was 15.32 years. Of the 300 children tested, 60 (20%) were infected. Urogenital schistosomiasis prevalence in males 38 (63%) was significantly higher ( $p < 0.05$ ) than in females 22 (36.7%). Of the six (6) schools visited, Government Secondary School Amenkawo had the highest prevalence rates of 18 (30%),  $p > 0.05$ . Among the different age groups, 11-15 years had the highest prevalence of 32 (53.3%), while significantly lower prevalence  $p < 0.05$  was

\*Corresponding author: E-mail: [joy\\_chikwendu@yahoo.com](mailto:joy_chikwendu@yahoo.com);

recorded in those 21 years and above 6 (10%). In addition to *Schistosoma* parasite, *Trichomonas vaginalis* was also recorded in children 16 years of age and above. Urinary schistosomiasis is a major public health problem affecting both males and females of different age groups and in different localities in Oju Local Government Area of Benue State, Nigeria.

**Keywords:** Urogenital schistosomiasis; trichomoniasis; school children; *Schistosoma* parasite.

## 1. INTRODUCTION

Schistosomiasis is a parasitic disease caused by digenetic trematodes of the genus *Schistosoma*. 90% of worldwide cases occur in sub-Saharan Africa [1]. More than 207 million people, 85% of who live in Africa, are infected with schistosomiasis and an estimated 700 million people are at risk of infection in 76 countries where the disease is considered endemic, as their agricultural work, domestic chores and recreational activities expose them to infested water [2]. The intermediate host for transmission of urinary schistosomiasis is snails of the genus *Bulinus* which is found in freshwater bodies. Transmission of the parasite takes place in permanent water bodies as well as in seasonal ponds or streams [3]. Although *Schistosoma haematobium* infection does not always result in clinical diseases as many infections are asymptomatic, schistosomiasis in children can cause anemia, stunted growth and a reduced ability to learn [4].

Prevalence rates of 14(6.4%) children was reported out of 218 school children before treatment with praziquantel tablets in Langai community, Mangu local government area, Plateau State [5]. High prevalence of 97 (58.1%) of urinary schistosomiasis among school children in Abeokuta has been reported [6]. High prevalence of 46.6% was also reported in Ogbadibo, Benue State, Nigeria [7]. Lower urinary schistosomiasis prevalence of 20.6% in males and 13.3% in females was reported among Hausa communities in Kano, Nigeria [8]. Younger people (school aged children) are most vulnerable to schistosomiasis infection and play an important role in its transmission; therefore school aged children should serve as a reference group for assessing the need for intervention in communities [9].

This research aimed to determine the difference in prevalence of the parasite between sexes, based on location and age in Oju Lga of Benue State as this could shed light on how to channel scarce control resources to maximize control outcomes in endemic communities.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

Across-sectional study was conducted in Oju LGA, Benue State, Nigeria. Oju is located in Southern Benue and comprises of 3 Zones (A, B and C).

### 2.2 Sampling Site

Six (6) schools consisting of Primary and Secondary Schools were sampled namely;

L.G.E.A Oburu, Government Secondary School Amenka-owo, Solid foundation Nursery/primary School Ega-Okpodom, Government Secondary School Ikachi, L.G.E.A Obusa, and Government Secondary School OdigoUwokwu.

### 2.3 Ethical Clearance

Consent was obtained from school authorities, parents, and assent was obtained from school children prior to sampling.

### 2.4 Specimen Collection

A total of 300 urine samples were collected between 10:00 am and 2:00 pm (10).

### 2.5 Laboratory Analysis of Samples

Following centrifugation, urine deposits were examined under a light microscope using x10 and x 40 objectives (10).

### 2.6 Statistical Analysis

Data obtained were analyzed using Chi-Square analysis.

## 3. RESULTS

Table 1 shows Prevalence of Urinary Schistosomiasis among school children in Oju L.G.A., Benue State. 50 urine samples were examined per school and the number that tested positive was recorded. Table 2 shows

Prevalence of *Schistosoma haematobium* infection among school children in relation to Age in Oju L.G.A. Benue State for both males and females Table 3 Shows organisms found in the urine samples of the participants. Apart from *S. haematobium*, other pathogens which were found in urine samples among the different age groups were recorded.

#### 4. DISCUSSION

Schistosomiasis remains a major public health problem in tropical and subtropical countries especially in areas with poor sanitary conditions and among people with limited knowledge of the disease [11], [12]. Our results revealed a high prevalence of 20% among school children in Oju Local Government Areas of Benue State. Similarly high prevalence of 38.6% and 45.4% was reported among primary and secondary school children in Buruku and Katsina-ala local government areas, Benue State [13].

There was significant difference in prevalence between males and females and in participants aged 21 years and above, prevalence in males was more than double the prevalence in females. This could be because in the Oju LGA, males tend to spend more time in the water bodies than females as male children tend to be very adventurous and spend long periods in water bodies thus increasing their chance of contact with *S. haematobium* cercariae and so, increasing their chances of infection. These results were in accordance with results by [14].

Prevalence rates differed significantly among age groups and was highest in children 11-15 years of age and least in students aged 21 years and above. This could be because as children grow older, they tend not to frequent streams for activities and the worms within them died naturally as *S. haematobium* adult worms have been reported to live on the average for 3-10 years [15]. The reduced prevalence observed in

**Table 1. Prevalence of Urinary Schistosomiasis among school children in Oju L.G.A., Benue State**

Schools	Samples examined	Number positive (%)
L.G.E.A. Oboru	50	12(20)
G.S.S. Amenka-owo	50	18(30)
Solid Foundation Nus/pri	50	4(6.7)
G.S.S. Ikachi	50	5 (8.3)
L.G.E.A. Obusa	50	11(18.3)
G.S.S. Odigo	50	10(16.7)
Total	300	60(20)

$p < 0.05$

**Table 2. Prevalence of *Schistosoma haematobium* infection among school children in relation to Age in Oju L.G.A. Benue State**

Age (Years)	Number examined	Number positive (%)	Male	Female
5-10	78	12 (20%)	8(13.3%)	4(6.7%)
11-15	80	32 (53.3%)	20 (33.3%)	12(20%)
16-20	73	10 (16.7%)	6 (10%)	4 (6.7%)
21 and above	69	6 (10%)	4 (6.7%)	2(3.3%)
Total	300	60(100%)	38(63%)	22(36.7%)

Significant at  $p < 0.05$ ; mean age of children 15.43 years

**Table 3. Organisms found in the urine samples of the participants**

Age group	Number examined	Pus cell (%)	RBC (%)	<i>T. vaginalis</i> (%)	Epithelia cell (%)
5-10	69	-	22(40.7%)	-	-
11-15	68	18 (31%)	24(41.4%)	-	8(13.8%)
16-20	86	16(19.8%)	12(14.8%)	3(3.7%)	14(17.3%)
≥21	77	10(15.4%)	8(12.3%)	6(9.2%)	8(12.3%)
Total	300	44(14.7%)	66(22%)	9(3%)	30(10%)

age group 21years and above could also be as a result of reduced egg secretion by adult worms with prolonged infection (chronic infection). For this reason, drug treatment for adults is advised in cases where adults grew up in highly endemic areas, presented with symptoms like haematuria when they were younger but as they grew older, became asymptomatic and are currently not shedding eggs in their urine; adult worms can sometimes survive as long as 40 years within humans [16].

Prevalence rates were highest in G.S.S. Amenka-owo, and lowest in Solid foundation nursery and primary school. Similar disparities in prevalence rates in communities within a study area were reported in Kwara State, Nigeria [17] by Abdulkareem et al. (2018).

From our research, other pathogens were present in the urine samples like pus cells, RBC, *Trichomonas vaginalis* and epithelial cells. Pus cells were significantly higher in those aged 11-15 years than in those aged 21 and above. While RBC were significantly higher in children aged 11-15 years and 5-10 years than in any other age group. Students aged 20 years and above had higher incidence of *T. vaginalis*. *T. vaginalis* in young adults could be indicative of sexual activities [18].

## 5. CONCLUSION

Urinary schistosomiasis is endemic in Oju LGA, Benue State, Nigeria.

Alternative water sources like boreholes and mass drug administration with praziquantel could go a long way in controlling the disease in the area.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. World Health Organization. Schistosomiasis. Fact sheet. 2017;115.
2. WHO. Schistosomiasis Fact Sheet No.115; 2010a. Available:<http://www.who.int/mediacentre/factsheets/fs115/en/index.html>.
3. Omonijo AO, Asaolu SO, Ofoezie IE. Schistosomiasis transmission and water contact pattern in River Ureje in Ado-ekiti Local Government Area, Ekiti State. J Parasitol. 2013;8(1):26-36.
4. WHO. Control of Urinary Schistosomiasis in School Children; 2012.
5. Banwat ME, Daboer JC, Envuladu EA, Lar LA, Ogbonna CC. Effect of mass chemotherapy with praziquantel on the presence of schistosomiasis in primary school children in Langai community of Plateau State. J. Trop. Med. 2011;13(2):119-23.
6. Ekpo UF, Laja-Deile A, Akinola SO, Sammy OS, Mafiana CF. Urinary schistosomiasis among preschool children in a rural community near abeokuta, Nigeria: Parasite Vector. 2010;3:58.
7. Mbata T, Orji M, Oguoma V. The prevalence of urinary schistosomiasis in ogbadibo local government area of Benue State, Nigeria. IJID. 2009;7.
8. Dawaki S, Al-Mekhlafi H, Ithoi I, Ibrahim J, Abdulsalam AM, Ahmed A, Sady H, Atroosh WM, Al-Areeqi MA, Elyana FN, Nasr NA, Surin J. Prevalence and risk factors of urinary schistosomiasis among Hausa communities in Kano State, Nigeria. Rev Inst Med Trop Sao Paulo. 2016;58:54.
9. Pereira AP, Favre TC, Galvao AF, Beck L, Barbosa CS, Pieri OS. The prevalence of schistosomiasis in school-aged children as an appropriate indicator of its prevalence in the community. Mem Inst Oswaldo Cruz. 2010;105(4):563-9.
10. Cheesbrough M. District laboratory practice in tropical countries. Part 2. Cambridge University Press, London. 2010;56.
11. Houweling TA, Karim-Kos HE, Kulik MC, Stolk WA, Haagsma JA, Lenk EJ, Richardus JH, de Vlas SJ. Socioeconomic inequalities in neglected tropical diseases: A systematic review. PLoS Negl Trop Dis. 2016;10:e0004546.
12. King CH. Parasites and poverty: The case of schistosomiasis. Acta Trop. 2010; 113:95-104.
13. Houmsou RS, Amuta EU, Sar TT. Profile of an epidemiological study of urinary schistosomiasis in two local government areas of Benue State, Nigeria. Int. J. Biomed. Res. 2012;1(1):39-48.
14. Chukwu DI, Okeke YO, Chukwu OOC, Ojemundia T, Agama CO, Chilom SC, Jemitola SO. Prevalence of *Schistosoma haematobium* infection among primary school pupils in Ganawuri village, Riyom

- L.G.A of Plateau State, Nigeria. IJPT. 2013;3(1):50-54.
15. Wilkins HA. The epidemiology of schistosome infections in man. In: The biology of schistosomes. Rollinson D, Simpson AJD, editors. London, UK: Academy Press. 1987;373-397.
  16. Colley DG, Bustinduy AL, Secor EW, King CH. Human schistosomiasis. Lancet. 2014; 383(9936):2253-64.
  17. Babamale OA, Kolawole OH, Abdulganiyu K, Abdulkareem OA, Ugbomoiko US. Urogenital schistosomiasis among school children and the associated risk factors in selected rural communities of Kwara State, Nigeria. J. Trop. Med. 2018; 6913918.
  18. Sutton M, Sternberg M, Koumans EH, McQuilan G, Berman S, Markowitz L. The prevalence of *Trichomonas vaginalis* infection among reproductive-age women in the United States, 2001-2004. Clin Infect Dis. 2007;45(10):1319-26.

---

© 2019 Chikwendu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/46858>