



The Physical Properties and Micronutrient Status of Mayo-gwoi Floodplain Soils, in Taraba State, Nigeria

P. K. Kefas^{1*}, M. E. Ukabiala² and C. V. Azuka³

¹Taraba State University, Jalingo, Nigeria.

²Kogi State College of Agriculture, Ochaja, Nigeria.

³Department of Soil Science, University of Nigeria, Nsukka, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2016/25777

Editor(s):

(1) Junhong Bai, School of Environment, Beijing Normal University, Beijing, China.

Reviewers:

(1) Ade Onanuga, Dalhousie University, Halifax, Canada.

(2) Alie Kamara, Njala University, Sierra Leone.

(3) Raimundo Jimenez Ballesta, Universidad Autonoma Madrid, Spain.

Complete Peer review History: <http://sciencedomain.org/review-history/14367>

Original Research Article

Received 19th March 2016

Accepted 16th April 2016

Published 27th April 2016

ABSTRACT

The aim of this research was to study the physical characteristics and the status of micronutrients in floodplain soils of Mayo-Gwoi in Taraba state, Nigeria. A detailed soil survey was conducted at the Mayo-Gwoi floodplain using rigid grid approach. Observations were made at 100m regular intervals. Two profile pits were dug and sampled from each of the three soil mapping units identified. The samples were analyzed and characterized as follows; texture vary from loamy to sandy clay loam, bulk density (1.30 g/cm³), particle density (2.63 g/cm³), water holding capacity (37.5%), water content at field capacity (36%) and wilting point (24%), zinc (1.0 Mg/Kg), iron (6.3 Mg/Kg), copper (0.5 Mg/Kg) and manganese (6.3 Mg/Kg). These soils showed some evidence of degradation and could be productive if subjected to appropriate management and maintenance.

Keywords: Soil survey; degradation; Mayo-Gwoi floodplain soils; rigid grid.

*Corresponding author: E-mail: pkefas@yahoo.com;

1. INTRODUCTION

Knowledge of soil physical and chemical properties are key to making agronomic and environmental decisions. However, the heterogeneous nature of these soil properties mainly due to changes in structural and chemical composition of soil minerals from microscale to global scale [1], makes this decision difficult. Soil physical properties such as texture, structure and porosity directly affect soil water movement and storage with subsequent effect on nutrient availability, plant water use and growth. The dynamics of soil moisture including the water retention capacity and soil nutrient status determine to a large extent the soil productivity. According to [2], micronutrients are as important in plant nutrition as macronutrients; though they simply occur in plants and soils in much lower concentrations. It has been observed that plants grown in micronutrient-deficient soils exhibit similar reductions in productivity as those grown in macronutrient-deficient soils. [3] have also reported that micronutrient deficiencies in crop plants are widespread worldwide. However, the deficiencies in these micronutrients often result to poor crop yields [4]. Micronutrients in soils exist in the form of elements in primary and secondary minerals; adsorbed to mineral and organic matter surfaces; incorporated in organic matter and microorganisms; incorporated into solution, depending on the source of the micronutrients.

Understanding the relationships and dynamics among these forms is essential for optimizing plant productivity in micronutrients deficient soils. The availability of micronutrients to plants is determined by both the total amount of the nutrient in the soil and the soil properties, crop species and variety, cropping systems, land use and soil management [5,6]. [7] also noted that the micronutrient availability in soils is a function of rate of replenishment from soil solids to soil solution. Thus, there is urgent need to assess the micronutrient status of Mayo Gwoi floodplain soils for better soil management and improved agricultural productivity.

It is of concern that in spite of the increasing interest in Fadama farming in Taraba State, there is no available literature on the hydro-physical properties and micronutrient levels of the floodplain soils. Thus, based on the realization that such information forms the background to an efficient and judicious use of the soil resources,

the objective of this study was to assess the physical properties and the status of micronutrients in the soils of Mayo-Gwoi floodplain.

2. MATERIALS AND METHODS

2.1 Description of the Study Area

The Mayo-Gwoi floodplain is located between latitudes 8° 53' and 8° 85' North and longitudes 11° 23' and 11° 75' East. It covers an area of about 120 ha. It's located in Jalingo city, the capital of Taraba State. The geology of the area has been described by Bawden [8] as a basement complex and the rocks are mainly pre-cambrian granitic and migmatite gneisses. Jalingo lies within the tropical hinterland climate region. The region is characterized by double maxima rainfall pattern which has about four months of dry season with relative humidity having generally over 80% in the morning and falls to between 50 and 79% in the afternoon. The dry and wet seasons are controlled by the annual migration of the intertropical zone of convergence (ITZC). The dry season is characterized by the dry dust, laden with harmattan winds coming across the Sahara desert and occurring between November and February of every year. The wet season sets in by April and lasts until October [9].

2.2 Sampling and Laboratory Analysis

Detailed soil survey was conducted at the Mayo-Gwoi floodplain soils which is the study site. A Rigid grid method was employed for surveying the area, baseline was drawn and traverses perpendicular to the baseline were cut and observations were made at 100 m regular intervals. Based on the information obtained in the field, three (3) mapping units were identified viz mapping units 1, 2 and 3 respectively. In mapping units 1 and 3, two profile pits each were sunk while mapping unit two had three profile pits making seven profile pits. Profile pits were sampled according to the pedogenetic horizons. Soil colour was determined using Munsell soil colour chart. Particle size distribution was determined by Bouyoucous hydrometer method using sodium hexametaphosphate (Calgon) as the dispersant and the textural class determination adopted was the USDA textural triangle method [10]. The bulk density was by core method [11]. The particle density was determined as

JALINGO TOPO. SHEET 215.




Fig. 1. Topographic map of Jalingo

described by [10]. Soil moisture content was determined using the gravimetric method as described by [10] while the water holding capacity of the soil was determined by the method of [12]. The porosity of the soil was determined according to the method described by [11] in [10].

3. RESULTS AND DISCUSSION

Soil textures were variable within the mapping units, surface texture ranged from loamy sand through sandy clay loam to clay. The sand contents of the profiles changed with depth, it has clay texture in both surface and subsurface

horizons, these variations may be due to differences in parent materials and topography [13]. The clay content of the soils was low while the sand content was high. Soil structures were also variable: being weakly developed in pedons that are moderately coarse-textured (pedons 1-5) and well developed in a pedon with high clay content. This lack of structural development in the horizons could be attributed to the effect of low water table [14]. The bulk density of the soils ranged from 1.22 g/cm³ to 1.44g/cm³ (Table 1). These values are considered safe for root penetration since root penetration may be hindered in soils having bulk density above 1.75 g/cm³ [15].

Table 1. Physical properties of Mayo-Gwoi flood plain soils

Mapping unit	Profile depth	B.D g/cm ³	PD g/cm ³	Porosity (%)	F.C (%)	WP (%)	AV H ₂ O (mm)	WHC (%)	Sand (%)	Silt (%)	Clay (%)	Textural class
MU1	0-18	1.42	2.65	46	25	14	11	37.1	56	21.2	22.8	Scl
	18-61	1.44	2.64	46	23	14	9	37.2	66	11.2	22.8	Scl
	61-82	1.41	2.64	47	25	16	9	37.2	62	13.2	26.8	Scl
	82-99	1.26	2.63	52	40	29	11	38.1	52.8	38	22.8	S
	99-144	1.35	2.62	45	22	14	8	32.2	54	5.2	40.8	Scl
	144-158	1.33	2.64	50	31	16	15	38.2	28	43.2	28.8	Cl
	158-178	1.24	2.64	53	37	28	9	32.2	47.2	40	52.8	Sl
	178-200	1.22	2.65	54	43	27	16	33.3	52.0	43.2	46.8	Sl
	0-14	1.32	2.63	50	32	23	9	40.2	10	61.2	28.8	Scl
	14-41	1.34	2.64	49	30	21	9	40.3	58	3.2	38.8	Sc
	41-78	1.49	2.64	44	21	13	8	40.2	58	19.2	22.8	Scl
	78-122	1.33	2.65	50	31	19	12	40.1	78	3.2	18.8	Sl
	122-141	1.4	2.63	47	26	15	11	39.2	74	3.2	22.8	Scl
	141-180	1.34	2.59	49	30	21	9	34.6	58	3.2	38.8	SC
180-200	1.42	2.62	46	24	14	10	35.7	52	5.2	42.8	Scl	
MU2	0-30	1.23	2.64	54	42	26	16	38.2	42	31.2	26.8	L
	30-50	1.29	2.65	51	34	15	18	38.4	8.0	29.2	62.8	C
	50-85	1.35	2.65	49	29	15	14	38.5	58	19.2	22.8	Scl
	85-140	1.32	2.59	50	32	18	14	33.3	35.2	58.0	26.8	L
	140-175	1.18	2.64	56	51	38	14	34.2	10	45.2	44.8	SL
	0-38	1.24	2.64	53	41	24	16	39.6	32	35.2	32.8	Cl
	38-62	1.33	2.64	50	31	19	12	33.4	10	47.2	42.8	Sc
62-120	1.24	2.65	53	40	23	17	40.6	42	23.2	34.8	Cl	
MU3	0-18	1.24	2.64	53	41	24	16	39.6	32	35.2	32.8	Cl
	18-54	1.22	2.64	54	42	26	17	39.4	58	21.2	20.8	Scl
	54-105	1.29	2.64	51	36	24	12	39.7	2.0	53.2	44.8	Sl
	105-134	1.22	2.63	54	42	26	17	35.6	8.0	51.2	40.8	Sc

Mapping unit	Profile depth	B.D g/cm ³	PD g/cm ³	Porosity (%)	F.C (%)	WP (%)	AV H ₂ O (mm)	WHC (%)	Sand (%)	Silt (%)	Clay (%)	Textural class
	134-184	1.44	2.64	46	24	13	10	34.7	4.0	19.2	22.8	Sc
	0-34	1.18	2.57	56	54	42	12	40.3	36	21.2	42.8	C
	34-52	1.22	2.61	54	46	34	12	40.5	22	5.2	72.8	C
	52-93	1.18	2.58	55	53	42	11	33.4	18	9.2	72.8	C
	93-130	1.25	2.63	53	41	31	10	35.4	40	3.2	56.8	C
	0-38	1.25	2.64	53	40	28	12	41.2	32	17.2	50.8	C
	38-60	1.24	2.61	53	43	32	11	40.6	34	9.2	56.8	C
	60-90	1.37	2.57	48	43	32	15	40.2	32	45.2	22.8	L
	90-120	1.25	2.64	53	41	29	12	40.4	32	16.8	51.2	C
	120-150	1.37	2.63	48	43	32	15	35.3	32	45.2	22.8	L
Range	-	1.18-1.49	2.57-2.65	44-56	21-54	13-42	8-18	32.2-41.2	2-78	3.2-61.2	18.8-72.8	
Mean	-	1.30	2.63	51	36	24	12	37.5	39	25.9	37.8	

BD= bulk density; PD =particle density; FC =field capacity; WP =wilting point; AV.H₂O=available water; WHC= water holding capacity, C=Clay, SC=Sandy clay, Cl=Clay loam, Scl=Sandy clay loam, L=Loam, S=Sand

Table 2. Micronutrients in soils of Mayo-Gwoi floodplain

Pedon number	Pedons depth (cm)	Zn (Mg/kg)	Fe (Mg/kg)	Cu (Mg/kg)	Mn (Mg/kg)
P1	0-18	1.1	9.7	ND	6.4
	18-61	2.6	8.1	0.5	4.4
	61-82	2.7	7.4	0.6	20.8
	82-99	2.1	5.6	ND	0.8
	99-144	2.6	3.1	0.5	4.6
	144-158	0.4	2.7	0.5	4.9
	158-178	0.5	2.8	0.5	10.2
	178-200	0.7	2.2	0.4	7.8
P2	0-14	1.5	8.1	0.5	5.9
	14-41	1.6	8.8	0.5	5.6
	41-78	1.6	5.4	0.5	9.2
	78-127	2.6	8.9	0.3	11.7
	127-141	2.4	5.6	0.4	3.1
	141-180	0.5	3.9	0.3	6.9
	180-200	0.3	7.7	0.7	4.0
P3	0-30	1.1	8.7	0.4	15.1
	30-50	0.5	6.9	0.4	6.3
	50-85	0.5	5.1	0.7	3.9
	85-140	0.8	7.8	0.5	7.1
	140-175	0.6	5.0	0.4	8.7
P3	0-32	0.5	6.2	0.6	2.4
	32-62	0.8	5.7	ND	6.1
	62-120	0.8	5.7	0.5	6.1
P4	0-18	0.3	5.6	0.3	2.8
	18-54	0.3	7.7	0.4	3.6
	54-105	0.5	6.0	0.4	6.9
	105-134	0.2	5.1	0.5	1.1
	134-184	0.6	5.0	0.6	7.2
P5	0-38	0.8	9.7	0.5	8.0
	38-52	0.7	5.9	0.5	6.3
	52-93	0.8	4.0	0.7	8.9
	93-130	0.6	7.1	0.4	9.2
P6	0-38	0.5	9.0	0.5	5.3
	38-60	0.6	5.7	0.6	5.4
	60-90	0.7	8.0	0.5	7.0
	90-120	0.7	8.7	ND	5.9
	120-150	0.1	3.4	0.3	6.4
Range	-	0.1-2.7	2.2-9.7	0.3-0.7	0.8-20.8
Mean	-	1.0	6.3	0.5	6.7

Note: ND = No data

The water holding capacity in both the surface and subsurface horizons of the soils were low (Table 2) and this could be attributed to the low organic matter content. Organic matter has been known for its importance at improving water retention capacity of most surface soils [13]. Consequently, water at field capacity, water at

wilting point and available water in the soils are low but fall within the range that cannot cause any negative effect to most arable crops [3]. The concentration or content of zinc in these soils were low (Table 2) and could be due to low organic matter content of the soils [13]. The iron content of the soils were generally low to

moderate (Table 2). The low iron content could be due to transformation and redox reactions [1]. Similar result was reported by [16] on the study of profile distribution of some Hydromorphic soils of Dass, Bauchi State. However, the level of available copper in the soils was deficient in all the pedons (Table 2) at both the surface and subsurface horizons which could be attributed to the low crystal concentration of copper in the soils dynamics [2]. The level of manganese was moderate at the surface and subsurface horizons in all the pedons (Table 2). This could also be attributed to the low organic matter content and the acidic nature of the soils shown by their pH values. [13] and [2] opined that the availability of most of the micronutrients in soils depend on the soil pH and organic carbon contents. Consequently, improving the organic carbon contents of the soil which serves as the mainstay of most extractable soil micronutrients [1], could help to improve the productivity of Mayo-Gwoi flood plain soils.

4. CONCLUSION

The study highlights the hydro-physical characteristics and the micronutrient status of soils from Mayo-Gwoi floodplain in Taraba state, Nigeria. The soils are characterized to have varying textures ranging from loam through loamy sand to sandy clay loam, low to moderate bulk density values as well as water holding capacity. The soils also showed low to moderate levels of the micronutrients (zinc, iron, copper and manganese). Good productivity of these soils would be ensured by adequate soil management strategies such as addition of organic matter in the form of poultry manure to improve the soil since organic matter is a major source of micronutrients in soils as well as improving soil physical health.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Brady NC, Weil RR. The nature and properties of soil (14th ed). Pearson Education Limited, USA; 2014.
2. Havlin SL, Nelson WL, Beaton JD, Tisdale JL. Soil fertility and fertilizers. 7th edition. Prentice-Hall of India. 2005;528.
3. Fageria NK, Baliger VC, Clark RB. Micronutrients in crop production. *Advances in Agronomy*. 2002;77:185-268.
4. Udo de Haes HA, Voortman RL, Bastein T, Bussink DW, Rougoor CW, vander Weijden WJ. Scarcity of micronutrients in soil, feed, food, and mineral reserves - Urgency and policy options. Platform for Agriculture, Innovation & Society; 2012.
5. Wei XR, Hao MD, Shao MG, Gale WJ. Changes in soil properties and the availability of soil micronutrients after 18 years of cropping and fertilization. *Soil and Tillage Res*. 2006;91:120-130.
6. Li BY, Huang SM, Wei MB, Zhang HL, Sheng AI, Xu JM, Ruan XL. Dynamics of soil and grain micronutrients as affected by long-term fertilization in an aquic inceptisol. *Pedosphere*. 2010;19:597-605.
7. Verma YK, Setia RK, Sharma PK, Singh Charanjit, Ashok Kumar. Pedospheric variations in distribution of DTPA-extractable micronutrients in soils developed on different physiographic units in central parts of Punjab, India. *International Journal of Agriculture and Biology*. 2005;7:243-246.
8. Bawden MG. The land systems, land resources study No. 9, LRD. Tolworth Towers Subritan Surrey, England. 1972; 205-209.
9. Iloeje NP. A new Geography of Nigeria. New Revised edition, Longman. Nig. Ltd. Ikeja. 1981;201.
10. Jaiswal PC. Soil, plant and water analysis. Kalyani Publishers, Ludhiana, New Delhi, NardaHyderabad, India. 2003;1-399.
11. Blake GR, Hartge KH. Bulk Density. In: methods of soil analysis, part 1. Physical and mineralogical methods. Klute A, (ed) American Society of Agronomy, Madison, WI USA. 1986;365-375.
12. Klute A. Water retention: Laboratory methods of soil analysis, part 1: Physical and mineralogical methods, 2nd ed. ASA, SSSA, Madison USA. 1986;635-660.
13. Brady NC, Weil RR. The nature and properties of soils. 13th edition. Pearson education. India; 2002.
14. Udo EJ. Nutrient status and agricultural potentials of wetland soils. A paper presented at the 27th Annual Conference of Soil Science Society of Nigeria, Calabar; 2001.
15. Esu IE. Characterization, classification and management problems of the major soil

- orders in Nigeria. 26th Inaugural Lecture of the University of Calabar. 2005;65.
16. Mustapha S, Udom GN, Umar AM. Profile distribution of some physico-chemical properties of some hydromorphic soils of Dass, Bauchi State, Nigeria. Nigeria Journal of Agriculture Technology. 2003;11:30-38.

© 2016 Kefas et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/14367>