

A Novel Dynamic Grey Action Quantity GM(1,1,b) Model and Its Application

Lang Yu^{1*}

¹School of Science, Southwest University of Science and Technology, Mianyang 621010, Sichuan, China.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/ARJOM/2019/v13i430113

Editor(s):

(1) Dr. Krasimir Yankov Yordzhev, Associate Professor, Faculty of Mathematics and Natural Sciences, South-West University, Blagoevgrad, Bulgaria.

Reviewers:

(1) Abdullah Sonmezoglu, Yozgat Bozok University, Turkey.
(2) Dr. Pasupuleti Venkata Siva Kumar, Vallurupalli Nageswara Rao Vignana Jyothi Institute of Engineering & Technology, India.

(3) Dr. Olumide Adesina, Olabisi Onabanjo University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49076>

Received: 01 March 2019

Accepted: 11 May 2019

Published: 21 May 2019

Method Article

Abstract

The classical GM(1,1) model treats the grey action quantity as an invariant constant, but changes have occurred within the system as time and space change. If the fixed grey action quantity is still used for modeling, the model will have errors. Aiming at this shortcoming, this paper proposes a GM(1,1,b) model in which the grey action quantity can be dynamically changed. Starting from the background value formula, the model solves the grey action quantity at different time points by the development coefficient, and fits the sequence with the DGM(1,1) model, then brings the obtained time response sequence into the classical GM (1, 1) to replaces the grey action quantity constant, so as to establish a GM(1,1,b) model with dynamic change of grey action quantity. Finally, the model is applied to the example of China's rural residents' consumption index. The numerical example shows that the GM(1,1,b) model proposed in this paper effectively improves the prediction accuracy of the model and verifies the effectiveness and practicability of the improved model.

Keywords: Rural residents' consumption level index; grey action quantity; GM(1,1,b); prediction accuracy.

1 Introduction

As one of the main driving forces for China's economic growth, consumption has affected the growth and speed of China's economy to a certain extent. Under the background of the continuous development of

*Corresponding author: E-mail: swust2018yl@163.com;

China's social economy, the consumption level of residents can effectively promote the growth of China's economy on the one hand, and the mutual influence with China's economic growth on the other. Both have a relationship of promotion and causality. At the same time, with the universal application of the Internet and the deepening of national policies in recent years, the quality of life of rural residents in China has been improved, and the level of consumption has also undergone significant changes. Therefore, scientific and reasonable prediction and analysis of China's rural residents' consumption level index can provide theoretical guidance for the macro-control of rural residents' consumption level.

In the prediction of the consumption level index, many scholars have conducted research. Among them, Meng [1] predicted the national consumption level index of China from 2010 to 2014 by establishing ARIMA model and residual autoregressive model. The results show that the ARIMA (2,2,0) model can achieve high prediction accuracy in short-term prediction. Li [2] used regression analysis to predict the consumption level of urban residents in Inner Mongolia, and applied computer to analyze the data. Wei [3] of Jilin university combined trend extrapolation method and exponential smoothing method, took the consumption level of urban residents in China from 1978 to 2007 as the analysis object, and comprehensively predicted the consumption level of urban residents by using deterministic time series analysis, and adopted the method of equal weight combination to improve the prediction accuracy. Ma et al. [4] improved a traditional PCA method and established a new evaluation model and applied it to the study of Qinghai urban residents' consumption level index, which can better evaluate the consumption level of residents. Li [5] used SAS software to construct ARIMA model for time series analysis and short-term forecast of rural residents' consumption level in China. The forecast results show that the consumption level of rural residents in China will be further improved rapidly, and then propose a development strategy that takes into account overall planning. Li et al. [6] used the GM (1,1) model and the buffer operator axioms, combined with qualitative analysis, to predict the development prospects of China's consumption level from 2000 to 2005. Cheng [7] believes that the ordinary grey prediction model can't accurately predict the consumption level index of rural residents in China, so an improved GM (1,1) model based on initial value correction is proposed to predict the consumption level index of rural residents in China in the future, and the model is validated with 2012 data.

Since the introduction of the grey system theory by the famous Chinese scholar Deng [8] in 1982, after more than 30 years of development, the grey prediction model has been widely used in economic, agricultural, military, ecological and other fields [9,10,11,12,13]. The grey prediction model has good performance in predicting "small sample" and "poor information" data sequences. It is an important part of the grey theory. The most classical prediction model is GM(1,1), where the first "1" represents the first order differential equation and the second "1" represents a variable. The improvement of the GM(1,1) model has never stopped, mainly in the following five aspects: (1) preprocessing of raw data [14,15,16]; (2) improvement of background value [17,18,19]; (3) initial value improvement [20]; (4) parameter improvement of the model [21,22]; (5) combination improvement of the above method [23].

In summary, there are not many literatures on the consumption index of rural residents, and there are no systematic methods and models. Starting from the classical GM (1,1) model, this paper proposes a novel GM (1,1) model with dynamic features. The discrete DGM(1,1) model is used to model and predict the grey action quantity sequence, which further improves the prediction accuracy of the GM(1,1) model and broadens the application range of the model. It provides a new modeling idea for the prediction of rural residents' consumption level index.

2 Basic Model

2.1 GM (1,1) model

Let $X^{(0)} = (x^{(0)}(1), x^{(0)}(2), \dots, x^{(0)}(n))$ as a non-negative raw sequence. Grey differential equations can be established for sequences satisfying smooth conditions. After a first order accumulation, $X^{(1)} = (x^{(1)}(1), x^{(1)}(2), \dots, x^{(1)}(n))$ is generated. $X^{(1)}$ is 1-AGO sequence of $X^{(0)}$, where

$$x^{(1)}(k) = \sum_{i=1}^k x^{(0)}(i), (k=1,2,\dots,n). \tag{1}$$

And $Z^{(1)} = (z^{(1)}(1), z^{(1)}(2), \dots, z^{(1)}(n))$ is the sequence mean generated of consecutive neighbors of $X^{(1)}$, where

$$z^{(1)}(k) = 0.5(x^{(1)}(k) + x^{(1)}(k-1)), k = 2, 3, \dots, n. \tag{2}$$

Thus, the grey differential equation is defined as

$$x^{(0)}(k) + az^{(1)}(k) = b, k = 2, 3, \dots, n, \tag{3}$$

where, a is the development coefficient and b is the grey action quantity. If $\hat{u} = (a, b)^T$ is a parameter sequence and there is

$$Y_1 = \begin{pmatrix} x^{(0)}(2) \\ x^{(0)}(3) \\ \vdots \\ x^{(0)}(n) \end{pmatrix}, B_1 = \begin{pmatrix} -z^{(1)}(2) & 1 \\ -z^{(1)}(3) & 1 \\ \vdots & \vdots \\ -z^{(1)}(n) & 1 \end{pmatrix}. \tag{4}$$

Then the least square estimate parameters sequence of the equation (3) satisfies

$$\hat{u} = (B_1^T B_1)^{-1} B_1^T Y_1. \tag{5}$$

If B_1, Y_1, \hat{u} satisfies equations (4) and (5), the corresponding whitening differential equation is

$$\frac{dx^{(1)}(t)}{dt} + ax^{(1)}(t) = b. \tag{6}$$

Let $x^{(0)}(1) = \hat{x}^{(0)}(1)$, get the time response sequence of the GM(1,1) model as

$$\hat{x}^{(1)}(k+1) = \left(x^{(0)}(1) - \frac{b}{a}\right)e^{-ak} + \frac{b}{a}, k = 1, 2, \dots, n-1, \dots. \tag{7}$$

After restoration, the predicted value $\hat{x}^{(0)}$ of $X^{(0)}$ is

$$\begin{aligned} \hat{x}^{(0)}(k+1) &= \hat{x}^{(1)}(k+1) - \hat{x}^{(1)}(k) \\ &= (1 - e^a) \left(x^{(0)}(1) - \frac{b}{a}\right) e^{-ak}, k = 1, 2, \dots, n-1, \dots. \end{aligned} \tag{8}$$

2.2 DGM (1,1) model

Let the non-negative sequence $X^{(0)}$ and the 1-AGO sequence $X^{(1)}$ as described in the above definition, called

$$\hat{x}^{(1)}(k+1) = \beta_1 \hat{x}^{(1)}(k) + \beta_2, \quad (9)$$

as a DGM (1,1) model, or as a discrete form of the GM (1,1) model [24]. The DGM(1,1) model is similar to GM(1,1), where the first "1" also represents the first order differential equation, and the second "1" also indicates that there is a variable.

If $\hat{\beta} = [\beta_1, \beta_2]^T$ is a sequence of parameters, and

$$Y_2 = \begin{pmatrix} x^{(1)}(2) \\ x^{(1)}(3) \\ \vdots \\ x^{(1)}(n) \end{pmatrix}, B_2 = \begin{pmatrix} x^{(1)}(1) & 1 \\ x^{(1)}(2) & 1 \\ \vdots & \vdots \\ x^{(1)}(n-1) & 1 \end{pmatrix}. \quad (10)$$

Then the least square estimate sequence of parameters $\hat{\beta} = [\beta_1, \beta_2]^T$ of the discrete grey prediction model

$\hat{x}^{(1)}(k+1) = \beta_1 \hat{x}^{(1)}(k) + \beta_2$ satisfies

$$\hat{\beta} = (B_2^T B_2)^{-1} B_2^T Y_2. \quad (11)$$

Let $\hat{x}^{(1)}(1) = x^{(0)}(1)$, get the recursive function as

$$\hat{x}^{(1)}(k+1) = \beta_1^k \left(x^{(0)}(1) - \frac{\beta_2}{1-\beta_1} \right) + \frac{\beta_2}{1-\beta_1}, k = 1, 2, \dots, n-1, \dots. \quad (12)$$

Restore value is

$$\begin{aligned} \hat{x}^{(0)}(k+1) &= \hat{x}^{(1)}(k+1) - \hat{x}^{(1)}(k) \\ &= (\beta_1 - 1) \left(x^{(0)}(1) - \frac{\beta_2}{1-\beta_1} \right) \beta_1^k, k = 1, 2, \dots, n-1, \dots. \end{aligned} \quad (13)$$

3 Dynamics of Grey Action Quantity b and Construction of GM(1,1,b) Model

Regarding the optimization and improvement of the GM(1,1) model, the optimization of the grey action quantity is rare. The grey action quantity is mined from the background value, which is exactly grey, which reflects the relationship of the data sequence. The traditional GM (1,1) model regards the grey action

quantity b as an invariant constant, that is, the external disturbance is regarded as constant, so the purpose of the processing is to facilitate the solution of the subsequent model and simplify the calculation of the model process. However, as can be seen from the literature [25,26,27,28,29,30,31], the amount of grey action quantity will change with time and space. If it is regarded as an invariant constant, this will inevitably lead to errors in the GM(1,1) model. The prediction accuracy of the model has a certain impact.

In general, when the original sequence $X^{(0)}$ changes from the first data $x^{(0)}(1)$ through time $t = 1, 2, \dots, n$ to the n th data $x^{(0)}(n)$, at this stage the environment of the entire system may have changed greatly with time. It can be seen from the grey differential equation $x^{(0)}(k) + az^{(1)}(k) = b, k = 2, 3, \dots, n$ of the classical GM(1,1) model that the classical GM(1,1) model lacks dynamic adjustment of the grey action quantity, which causes the model to not dynamically adapt to changes in the system over time. This is also one of the key reasons for the low accuracy of GM(1,1) model fitting and predicting.

Starting from the grey differential equation of the GM(1,1) model, when $k = 2, 3, \dots, n$ is considered, the grey action quantity of GM(1,1) model are respectively

$$k = 2, b^{(0)}(1) = x^{(0)}(2) + az^{(1)}(2), \tag{14}$$

$$k = 3, b^{(0)}(2) = x^{(0)}(3) + az^{(1)}(3), \tag{15}$$

⋮

$$k = n, b^{(0)}(n-1) = x^{(0)}(n) + az^{(1)}(n). \tag{16}$$

The grey action quantity sequence $B = (b^{(0)}(1), b^{(0)}(2), \dots, b^{(0)}(n-1))$ can be obtained by the above formulas (14), (15), and (16). Where, a is the development coefficient of the GM(1,1) model, and $z^{(1)}(k)$ is the sequence mean generated of consecutive neighbors of $X^{(1)}$.

Since the grey action quantity sequence $B = (b^{(0)}(1), b^{(0)}(2), \dots, b^{(0)}(n-1))$ is also a time series, so the grey action quantity sequence $B = (b^{(0)}(1), b^{(0)}(2), \dots, b^{(0)}(n-1))$ is used as the raw data sequence to establish a discrete DGM(1,1) model for fitting the grey action quantity sequence and predicting the change trend of the grey action quantity. Let $\hat{b}^{(1)}(1) = b^{(0)}(1)$, then the recursive function is

$$\hat{b}^{(1)}(k+1) = \beta_1^k \left(b^{(0)}(1) - \frac{\beta_2}{1-\beta_1} \right) + \frac{\beta_2}{1-\beta_1}, k = 1, 2, \dots, n-2, \dots. \tag{17}$$

Restore value is

$$\begin{aligned} \hat{b}^{(0)}(k+1) &= \hat{b}^{(1)}(k+1) - \hat{b}^{(1)}(k) \\ &= (\beta_1 - 1) \left(b^{(0)}(1) - \frac{\beta_2}{1-\beta_1} \right) \beta_1^k, k = 1, 2, \dots, n-2, \dots. \end{aligned} \tag{18}$$

The fitted values obtained by the above formula (18) and the predictive values $\hat{b}^{(0)}(k)$ are brought into the above equations (7) and (8) to achieve dynamic change of the grey action quantity.

$$\hat{x}^{(0)}(k+1) = (1 - e^a) \left(x^{(0)}(1) - \frac{(\beta_1 - 1) \left(b^{(0)}(1) - \frac{\beta_2}{1 - \beta_1} \right) \beta_1^k}{a} \right) e^{-ak}, \quad (19)$$

$$k = 1, 2, \dots, n, \dots$$

The above formula (19) is a GM(1, 1, b) model in which the grey action quantity obtained by combining the GM(1, 1) model and the DGM(1, 1) model can be dynamically changed. The model implements a dynamic change of grey action quantity that allows the model to adapt to changes in the external environment. In theory, the model should have better prediction accuracy than the static grey action quantity.

4 Example Application and Analysis

In order to verify the validity and practicability of GM(1,1,b) model established in this paper. The consumption level index of rural residents' in China from 2002 to 2013 was obtained from the Statistical Yearbook of China, and used as an example to analyze the data for modeling and prediction analysis. Compare the fitted and predicted accuracy of the proposed GM(1,1,b) improved model and the classical GM(1,1) model. The raw data is shown in the following Table 1.

Table 1. China's 2002-2013 rural residents consumption level index (1978=100)

Year	2002	2003	2004	2005	2006	2007
	421.10	440.50	457.80	488.90	524.70	570.40
Year	2008	2009	2010	2011	2012	2013
	610.30	666.90	716.00	808.60	880.40	955.80

In this paper, the consumption level index of rural residents from 2002 to 2008 was selected as the fitting modeling data, and the consumption level index of rural residents from 2009 to 2013 was used as the predictive comparison data to establish the classical GM(1,1) model and GM(1,1,b) model respectively. After calculation, the estimated parameter of GM(1,1) model is

$$\hat{u} = (\hat{a}, \hat{b})^T = (-0.0685, 387.7236)^T. \quad (20)$$

From the above estimated parameter column \hat{u} , the time response sequence of the GM(1,1) model can be obtained as

$$\hat{x}^{(0)}(k+1) = (1 - e^{-0.0685}) \left(x^{(0)}(1) + \frac{387.7236}{0.0685} \right) e^{0.0685k}, \quad (21)$$

$$k = 1, 2, \dots, n, \dots$$

The grey action quantity sequence

$$B = [396.59, 381.1, 383.51, 385.93, 388.38, 390.83, 393.31, 395.8, 398.3, 400.82, 403.36],$$

can be calculated by the development coefficient a , and the DGM(1,1) model of the grey action quantity amount is established by using the sequence to fit and predict the grey action quantity amount. After calculation, the estimated parameters of the DGM(1,1) model is

$$\hat{\beta}=(\beta_1, \beta_2)^T=(1.0063, 378.585)^T . \quad (22)$$

From the estimated parameter column $\hat{\beta}$ of the above DGM(1,1) model, the DGM(1,1) model recursive function can be calculated as

$$\hat{b}^{(1)}(k+1)=1.0063^k\left(b^{(0)}(1)+\frac{378.585}{0.0063}\right)-\frac{378.585}{0.0063}, \quad (23)$$

$$k=1, 2, \dots, n-2, \dots .$$

Restore value is

$$\hat{b}^{(0)}(k+1)=0.0063\left(b^{(0)}(1)+\frac{378.585}{0.0063}\right)1.0063^k, \quad (24)$$

$$k=1, 2, \dots, n-2, \dots .$$

Bring the equation (24) into equation (21) to obtain the GM (1,1,b) model time response sequence as

$$\hat{x}^{(0)}(k+1)=\left(1-e^{-0.0685}\right)\left(x^{(0)}(1)+\frac{0.0063\left(b^{(0)}(2)+\frac{378.585}{0.0063}\right)1.0063^k}{0.0685}\right)e^{0.0685k}, \quad (25)$$

$$k=1, 2, \dots, n, \dots .$$

In order to compare and analyze the GM (1,1) model and the GM (1,1,b) model, $MPAE_{pred}$ and $MAPE_{tol}$ are used as evaluation indicators of the model. Where $MPAE_{pred}$ is the mean absolute percentage error of the extrapolated predicted value and $MAPE_{tol}$ is the total mean absolute percentage error. The calculation formula are as follows

$$MAPE_{tol}=\frac{1}{n} \sum_{k=1}^n\left|\frac{x^{(0)}(k)-\hat{x}^{(0)}(k)}{x^{(0)}(k)}\right| \times 100\%, \quad (26)$$

$$MPAE_{pred}=\frac{1}{n-N} \sum_{k=N+1}^n\left|\frac{x^{(0)}(k)-\hat{x}^{(0)}(k)}{x^{(0)}(k)}\right| \times 100\%. \quad (27)$$

Where n represents the total number of sample data and N represents the number of data used to fit the model.

The fitted and raw data and relative errors of the GM(1,1) and GM(1,1,b) models are shown in Table 2. The five-steps extrapolated prediction and prediction errors for the two models are shown in Table 3.

Table 2. GM(1,1) and GM(1,1,b) model fitted and relative errors comparison (%)

Year	Raw data	GM(1,1)	Relative errors	GM(1,1,b)	Relative errors
2002	421.10	421.10	0.00	421.10	0.00
2003	440.50	431.14	2.12	440.32	0.04
2004	457.80	461.69	0.85	438.30	4.26
2005	488.90	494.40	1.13	494.57	1.16
2006	524.70	529.43	0.90	535.24	2.01
2007	570.40	566.95	0.61	579.07	1.52
2008	610.30	607.12	0.52	626.31	2.62

Table 3. Comparison of GM(1,1) and GM(1,1,b) model prediction and prediction errors (%)

Year	Raw data	GM(1,1)	Relative errors	GM(1,1,b)	Relative errors
2009	666.90	650.14	2.51	677.21	1.55
2010	716.00	696.20	2.77	732.05	2.24
2011	808.60	745.53	7.80	791.15	2.16
2012	880.40	798.36	9.32	854.83	2.90
2013	955.80	854.93	10.55	923.44	3.39
$MPAE_{pred}$		6.5902		2.4471	
$MAPE_{tol}$		3.2566		1.9873	

It can be seen from Table 3, $MPAE_{pred}$ and $MAPE_{tol}$ of the GM (1,1,b) model are only 2.4471% and 1.9873%, while $MPAE_{pred}$ and $MAPE_{tol}$ of the GM (1,1) model are as high as 6.5902% and 3.2566%, which is obviously higher than the GM (1,1,b) model proposed in this paper.

Fig. 1 visually compares the fitted and predictive values of the GM (1,1) and GM (1,1,b) models, and Fig. 2 compares the prediction errors of the GM (1,1) and GM (1,1,b) models. The above two figures clearly show that the GM(1,1,b) model proposed in this paper is superior to the classical GM(1,1) model.




Fig. 1. Comparison of fitted and predicted values of GM(1,1) and GM(1,1,b) models




Fig. 2. Prediction errors of GM(1,1) and GM(1,1,b) models

5 Conclusion

This paper starts from the classical GM(1,1) model, aiming at the shortcoming that GM(1,1) model ignores the change of external environment and regards grey action quantity as a constant. By studying the dynamic change property of grey action quantity, the DGM(1,1) model was used to simulate and predict the grey action quantity. Based on this, a novel GM(1,1,b) model with dynamic grey action quantity can be proposed and applied to the prediction of rural residents' consumption level index. The results show that the proposed GM(1,1,b) model has higher prediction accuracy than the classical GM(1,1) model, which verifies the practicability and effectiveness of the GM(1,1,b) model. At the same time, it also provides relevant theoretical basis for the state to adjust the consumption index of rural residents in China.

Acknowledgements

This paper has been supported by Undergraduate Innovation Fund Project Accurate Funding Special Project by Southwest University of Science and Technology (No. JZ19-057).

Competing Interests

Author has declared that no competing interests exist.

References

- [1] Meng YB. Application of time series analysis in forecasting consumption level index. Journal of Guangxi Teachers Education University (Natural Science Edition). 2012;29(01):37-44.
- [2] Li ZY. Using regression analysis to predict the consumption level of urban residents in our district. Journal of Inner Mongolia University for Nationalities. 2008;14(06):73-74.

- [3] Wei WK. Analysis and prediction of consumption level of urban residents in china-analysis of combination model based on trend extrapolation and exponential smoothing method. *Prices Monthly*. 2011;03:74-77.
- [4] MA ZM, Geng SL, Shang C. Resident consumption level assessment model and its application based on multivariate statistical analysis. *Mathematics in Practice and Theory*. 2009;39(20):48-52.
- [5] Yan JB, Li Q. Forecast and analysis of rural households consumption level in china based on the ARIMA model. *Journal of Anhui Agricultural Sciences*. 2011;39(14):8754-8756.
- [6] Li SW, Wang AR. The grey calculation of households consumption. *Journal of Gansu Sciences*, 2002;03:100-104.
- [7] Cheng YL, Wang B. Predictive analysis of china's rural residents' consumption level index-an improved GM(1,1) model based on initial value correction. *Rural Economy and Science-Technology*. 2014;25(08):113 -115.
- [8] Deng JL. Control problems of grey systems. *Systems & Control Letters*. 1982;1(5):288-294.
- [9] Fei ZH, Jiang ZY, He ZF. Discrete GM(1,1) model and its application for forecasting of real estate prices. *International Conference on Management Science & Industrial Engineering*; 2011.
- [10] Truong DQ, Ahn KK. An accurate signal estimator using a novel smart adaptive grey model SAGM(1,1). *Expert Systems with Applications*. 2012;39(9):7611-7620.
- [11] Li JL, Liao RQ, Chen XC. Generalized accumulated GM(1,1) cosine model and its application *IEEE International Conference on Grey Systems & Intelligent Services*; 2011.
- [12] Ou SL. Forecasting agricultural output with an improved grey forecasting model based on the genetic algorithm. *Computers & Electronics in Agriculture*. 2012;85(85):33-39.
- [13] Hsu LC, Wang CH. Forecasting integrated circuit output using multivariate grey model and grey relational analysis. *Expert Systems with Applications*. 2009;36(2):1403-1409.
- [14] Zeng B, Duan HM, Bai Y, Meng W. Forecasting the output of shale gas in China using an unbiased grey model and weakening buffer operator. *Energy*. 2018;151:238-249.
- [15] Liao RJ, Bian JP, Yang LJ, Grzybowski S, Wang YY, Li J. Forecasting dissolved gases content in power transformer oil based on weakening buffer operator and least square support vector machine-Markov. *IET Generation, Transmission & Distribution*. 2012;6(2):142-151.
- [16] Truong DQ, Ahn KK. An accurate signal estimator using a novel smart adaptive grey model SAGM(1,1). *Expert Systems with Applications*. 2012;39(9):7611-7620.
- [17] Wang ZX, Dang YG, Liu SF, Zhou J. The optimization of background value in GM(1,1) model. *Journal of Grey System*. 2007;10(2):69-74.
- [18] Wang ZX, Dang YG, Liu SF. Optimization of background value in GM(1,1) model. *Systems Engineering-theory & Practice*. 2008;28(2):61-67.
- [19] Wang YH, Liu Q, Tang JR, Cao WB, Li XZ. Optimization approach of background value and initial item for improving prediction precision of GM(1,1) model. *Journal of Systems Engineering and Electronics*. 2014;25(1):77-82.

- [20] Chen QM, Li J. Research on optimum weighted combination GM(1,1) model with different initial value. International Conference on Intelligent Computing. Springer, Cham. 2015;354-362.
- [21] Xie NM, Liu SF. Novel methods on comparing grey numbers. Applied Mathematical Modelling, 2010;34(2):415-423.
- [22] Zeng B, Liu SF, Xie NM. Prediction model of interval grey number based on DGM(1,1). Journal of Systems Engineering and Electronics. 2010;21(4):598-603.
- [23] Wang YH, Liu Q, Tang JR, Cao WB, Li XZ. Optimization approach of background value and initial item for improving prediction precision of GM(1,1) model. Journal of Systems Engineering and Electronics. 2014;25(1):77-82.
- [24] Zeng B, Liu SF. DGM(1,1) direct modeling approach of DGM(1,1) with approximate non-homogeneous exponential sequence. Systems Engineering —Theory & Practice. 2011;31(02):297-301.
- [25] Xiao XP, Wang HH. Change of GM(1,1, α) model background value on the influences of relative errors. Systems Engineering —Theory & Practice. 2014;34(02):408-415.
- [26] Xu HF, Liu SF, Fang ZG. Optimum grey action quantity for GM(1,1). Mathematics in Practice and Theory. 2010;40(02):26-32.
- [27] Xu ZD, Liu FX, Liu J. Optimum grey action quantity for GM(1,1) with fractional order accumulate and its research. Mathematics in Practice and Theory. 2016;46(19):235-242.
- [28] Liu CL, Qu SJ, Liu SQ, Chen J. Optimum whitenization equation for GM(1,1) model. Mathematics in Practice and Theory. 2015;45(04):213-220.
- [29] Wang J. Optimization and application of a new type of grey prediction model. Statistics & Decision. 2015;17:86-88.
- [30] Liu CL, Chen J, Qian JJ, Sun YL, Han XM. Optimum grey action quantity for GM(2,1) model. Mathematics in Practice and Theory. 2017;47(18):177-184.
- [31] Huang H. A new grey prediction model for dynamic development coefficient a. Statistics and Decision. 2016;21:19-21.

© 2019 Yu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<http://www.sdiarticle3.com/review-history/49076>