



Utilising Inter-Random Amplified Polymorphic DNA, Assess Genetic Variability in Indigenous Medicinal Plant Heart-leaved Moonseed (*Tinospora cordifolia*)

B. C. Akhilraj^{a++*}, J. Suresh^{b#}, K. Rajamani^{c#}, M. Kumar^{dt}
and R. Gnanam^{e#}

^a Department of Plantation, Spices, Medicinal and Aromatic Crops, HC & RI, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

^b Coconut Research Station, Tamil Nadu Agricultural University, Aliyar Nagar, Pollachi, Tamil Nadu, India.

^c Department of Floriculture and Landscape Architecture, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

^d ICAR- Krishi Vigyan Kendra, Tamil Nadu Agricultural University, Tindivanam, Villupuram, Tamil Nadu, India.

^e Department of Bio Informatics, CPMB & B, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i132995

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/99494>

Original Research Article

Received: 08/03/2023

Accepted: 10/05/2023

Published: 19/05/2023

⁺⁺ Ph.D Scholar;

[#] Professor and Head;

[†] Professor (PB&G) & Programme Coordinator;

*Corresponding author: E-mail: bcakhilraj@gmail.com;

ABSTRACT

The Ayurvedic medical system, which uses herbal preparations for its traditional remedies, has been used in India since ancient times. One of the powerful immunomodulators is *Tinospora cordifolia*, also known as AMRITA, which is credited to this miracle medication for giving its user youth, longevity, and vigour. RAPD has been widely and successfully used to identify several species of medicinally significant plant. The need for research into and protection of therapeutic plant species is growing. The perseverance of the evaluation was to assess the genetic diversity among 10 accessions of *T. cordifolia* obtained from various districts in Tamil Nadu. Eighteen decamer primers were used in the study. Nine of them had repeatable, different polymorphism patterns in their primers. The DNA recovered from several *T. cordifolia* accessions exhibited random amplification and various levels of genetic polymorphism was observed. In the sample from Salem district the most genetic polymorphism was found.

Keywords: *Tinospora cordifolia*; Tamil Nadu; RAPD; genetic diversity; medicinal plant.

1. INTRODUCTION

Since ancient times, people have employed plants as medicine. The Ayurvedic medical system, which uses herbal preparations for its traditional remedies, has been used in India since ancient times. The therapeutic benefits of these medicinal plants have been thoroughly investigated in a number of scientific fields, including pharmacology, cosmetics, fragrance, cuisine, etc [1]. The term "Return to Nature" can be used to describe the recent shift in Indian medicine from allopathic to herbal therapy. Since millennia, medicinal plants have been highly valued for their curative qualities in treating ailments and preventing disease. Plants have the capacity for both natural healing and acting as potential antibacterial agents. According to estimations from the World Health Organization (WHO), 80% of people worldwide depend on traditional ayurvedic medicines to maintain a healthy way of living [2].

T. cordifolia is also referred as "Guduchi" [3], although it is most often referred to as "Giloya" a mythological Hindu term that refers to the holy concoction that prevents the godly spirits from ageing and keeps them youthful eternally. *T. cordifolia*, a herbaceous vine that grows 300 metres above sea level and found throughout the tropical Indian subcontinent and China, is a member of the Menispermaceae family. Although *T. cordifolia* is a resilient plant that can be cultivated in virtually any temperature, it loves a warm environment for the best development. The greatest time to grow this plant is during the rainy season, which is from July to August. This medicinal plant has the amazing capacity to generate aerial roots that may extend 10 metres down and 10 metres up from the earth. These

aerial roots are crucial to the plant's ability to grow again. This plant doesn't actually die; instead, it either drops to the ground or expands to a new "life line," reinstating itself. *T. cordifolia* is a climber that needs help from the outside to flourish. Neem is known as NEEM GILOY when it helps *T. cordifolia* flourish. Giloy and neem share a similar chemical makeup and exhibit superior medicinal qualities when combined [4].

T. cordifolia has a number of documented medical applications in Ayurveda. According to "Ayurveda," the nutrient-rich starch extracted from *T. cordifolia*'s roots and stems can treat diarrhoea and dysentery. Ayurvedic "Rasayanas" derived from *T. cordifolia* strengthen the immune system and increase the body's resistance to numerous illnesses [5]. The fresh plant components are thought to be more effective at treating illnesses. The plant's aqueous extract, known as Indian quinine, is effective in lowering high body temperatures brought on by a cold or dyspepsia. Along with these other uses, it is widely used for spermatorrhea, impotence, sclerotic disorders, urinary tract infections, acidosis, simple debility, and sexually transmitted illnesses. In addition to being used to cure leprosy and malaria, *T. cordifolia* roots also serve as natural sedatives [6].

A contemporary polymerase chain reaction-based approach for identifying genetic changes across and within the same species is called random amplified polymorphic DNA (RAPD). The DNA sequences of RAPD markers are amplified using a single, short, and randomly chosen oligonucleotide primer (usually 10 bases long). RAPD has been widely and successfully used to identify several species of medicinally significant

plant [7]. The need for research into and protection of therapeutic plant species is growing [8]. The following goal have been set for the current study by taking into account the significance of *T. cordifolia* as a medicinal plant and RAPD as a potent technique to evaluate the genetic polymorphism of medicinal plants.

2. MATERIALS AND METHODS

Plant collection: *Tinospora cordifolia* leaves were collected in Tamil Nadu, India, from a variety of biological zones and altitudes ranging from 8 to 378 m MSL. Entire plant samples were carefully categorised and brought back from the fields in ice boxes. The current research includes ten accessions of *T. cordifolia* from Trichy, Dindigul, Ariyalur, Ranipet, Vellore, Cuddalore, Coimbatore, Nagapattinam, Dindigul and Thirupathur (Table 1). Altitude was used as a reference point among other factors while examining genetic variation.

Sample collection: *T. cordifolia* stems were gathered in ten different places in Tamil Nadu, ranging from 8.00 to 378.00 meters above MSL (Table 1) over the years of 2022. Plant samples were collected from Thittacheri village of District Nagapattinam at the peak of 8m above MSL (Mean Sea Level), one town from municipality Cuddalore i.e Virudhachalam at the peak of 33m above MSL one town from municipality Ariyalur i.e Jayankondam at the peak of 61m above MSL one town from municipality Ranipet i.e Kalavai at the peak of 132m above MSL, one town from municipality Salem i.e Thalaivasal at the peak of 162m above MSL, one town from municipality Dindigul i.e Vedasandur at the peak of 212 m above MSL, one village from District Vellore i.e Gudiyatham at the peak of 268m above MSL, one town from municipality Coimbatore i.e Kinethukadavu at the peak of 308m above MSL, one village from District Thirupathur i.e Vaniyampadi at the peak of 341m above MSL, and one village from District Dindigul i.e Palani at the peak of 378m above MSL. The RAPD analysis and antibacterial properties of *T. cordifolia* have both been the subject of several investigations. For instance, Mohapatra and Rout, [9] reported fresh plant samples were gathered from Kanyakumari District Tamil Nadu, while leaves of *T. cordifolia* were taken from

Ambasamudram, Tirunelvi, Sankarankoil, Tenkasi and Alangulam, in District Tamil Nadu [10].

Extraction of genomic DNA using the modified CTAB technique: *Tinospora cordifolia* leaves yielded DNA (5gms). Using a pre-cooled mortar and pestle, leaves were pounded into a fine powder and then dissolved in liquid nitrogen. The powder was added to a centrifuge tube with 7 ml of preheated (65^o C) DNA extraction buffer, and the mixture was forcefully agitated to create a homogeneous suspension. The suspension was incubated at 65^o C for an hour, with inversion mixing performed every ten minutes. The mixture was extracted for around 15 minutes using an equal amount of phenol and chloroform (24:1). After incubation, centrifuged the samples at 10,000 rpm for 10 minutes. Transferring the top aqueous layer with care to a different centrifuge tube 0.6 litre of cold ethanol was added to this and stirred by inverting. Overnight, the samples were incubated at -20°C. The next morning, the samples were centrifuged for 10 minutes at 10,000 rpm after being mixed by inverting them. The particle was washed three times with 70% ethanol after the supernatant was discarded. The pellet was dried by air overnight. After drying, the pellet was dissolved the in 50 µl of TE [11].

PCR-RAPD- amplification: The Gene Amp PCR System 9700 R (Applied Biosystems, Carlsbad, USA) was used for the PCR amplification, with the following amplification parameters: initial denaturation at 94°C for 5 min, followed by 45 cycles of denaturation at 94°C for 1 min, primer annealing at 37°C for 1 min, amplification at 72°C for 2 min, and a final extension at 72°C for 5 min. Amplified DNA was electrophoresed in a 1.4% agarose gel using a 1µ TAE buffer electrophoresis at 50 V for three hours Ethidium bromide was used to observe the electrophoresis gel. Gene Ruler™, a 1-kb DNA ladder, was applied as usual. Eighteen primers were used for polymorphism analysis (Table 2).

The presence or lack of bands was used to assess genetic similarities. DNA bands may be seen and rated as present (1), absent (0), or not present, and a binary matrix was created. MVSP software was used for data processing, and genetic diversity was investigated.

Table 1. Samples from various Tamil Nadu locations collected for RAPD analysis

S. No.	Place	District	Latitude	Longitude	Altitude (MSL)	Code
1.	Thittacheri	Nagapattinam	10 ^o 52'09"N	79 ^o 47'08"E	8 m	TCNT
2.	Virudhachalam	Cuddalore	11 ^o 30'52"N	79 ^o 19'31"E	33 m	TCCV
3.	Jayankondam	Ariyalur	11 ^o 12'45"N	79 ^o 21'45"E	61 m	TCAJ
4.	Kalavai	Ranipet	12 ^o 46'01"N	79 ^o 25'01"E	132 m	TCRK
5.	Thalaivasal	Salem	11035'10"N	78045'30"E	162 m	TCST
6.	Vedasandur	Dindigul	10 ^o 31'50"N	77 ^o 56'54"E	212 m	TCDV
7.	Gudiyatham	Vellore	12 ^o 56'12"N	78 ^o 52'46"E	268 m	TCVG
8.	Kinethukadavu	Coimbatore	10 ^o 48'39"N	77 ^o 01'20"E	308 m	TCCK
9.	Vaniyampadi	Thirupathur	12 ^o 41'23"N	78 ^o 37'09"E	341 m	TCTV
10.	Palani	Dindigul	10 ^o 26'57"N	77 ^o 30'58"E	378 m	TCDP

Table 2. Primers used in the study for genetic variability of 10 *Tinospora cordifolia*

S. No.	Codes	SEQUENCE	Tm ^o c
1.	OPA - 01	5' GA GT CT CA GG 3'	32.00
2.	OPC - 02	5' GG TC TA GA GG 3'	32.00
3.	OPC - 14	5' GG AG TA CT GG 3'	32.00
4.	OPA - 03	5' AA GC CT CG TC 3'	32.00
5.	OPA - 09	5' GT CC CG AC GA 3'	32.00
6.	OPC - 04	5' AG CC AG CG AA 3'	32.00
7.	OPC - 15	5' TC TG TG CT GG 3'	32.00
8.	OPA - 13	5' GA CC GC CT TG T 3'	32.00
9.	OPA - 17	5' GG GT AA CG CC 3'	32.00
10.	OPA - 20	5' GT TT GC GA TC C 3'	32.00
11.	OPG - 11	5' GT GA GA CG GA 3'	32.00
12.	OPC - 11	5' AC CG AG CG AT 3'	32.00
13.	OPG - 02	5' TG TG TG CT AA 3'	32.00
14.	OPA - 04	5' GA TT GC CA AG T 3'	32.00
15.	TC - 01	5' GG CT AA CG GC 3'	32.00
16.	TC - 02	5' GG AC TG CA GA 3'	32.00
17.	TC - 03	5' AG TC AG CC AC 3'	32.00
18.	OPG - 24	5' CA GG CC CT TC 3'	32.00

3. RESULTS AND DISCUSSION

Researchers have identified 152 species in India that might be used to create ayurvedic medications, and *T. cordifolia* is one of these potentially useful plants [12]. RAPD markers have the capacity to detect and define the genetic diversity of plants [13]. Proper plant management of its genetic resources is aided by the many genetic components found in the crop and detected by markers reported Saha and co-workers [14]. Despite the availability of several molecular markers, RAPD markers are explored in the current research because of their implications.


***Tinospora cordifolia* molecular profiling with RAPD markers:** The genetic link and degree of relatedness among the 10 *T. cordifolia* accessions were discovered using RAPD primers. Eight primers were chosen from among them based on the highest level of polymorphism found in all DNA samples extracted from *Tinospora* leaves collected from various places in Tamil Nadu. The resultant amplified fragments

were examined for genetic variants and relationships based on the bands are present or not. DNA bands were either present means indicted as one (1) or absent means indicted as zero (0), depending on data. After RAPD, the bands were seen under a UV transilluminator, and gel documentation system photographs were made, as shown in Fig. 1 (I-VIII). All the analysed samples revealed polymorphism. Less polymorphic primers included OPC – 15, OPC - 04, OPA -13 and OPA - 09 whereas markers OPA – 03 and OPC - 14 had fair to outstanding polymorphism, as did primers OPA-01 and OPC-2. Table 3 provides an overview of primer codes, number of amplified alleles, number of polymorphic alleles, number of monomorphic alleles, polymorphic per cent and amplification status.

Initial RAPD results showed that 8 RAPD primers, namely OPC 2, OPA 01, OPC 14, OPA 13, OPA 03, OPA 09, OPC 1 and OPC 4 with 10 distinct samples of *T. cordifolia* accessions obtained from several districts in Tamil Nadu,

exhibited amplification and varying degree of genetic polymorphism. In the sample taken from Thalaivasal in the District of Salem, primer OPA 01 demonstrated good polymorphism, while samples taken from other locations revealed significant genetic variety. Fig. 1-I displayed all of the distinct bands produced by amplifying using OPA 01 primer. Using Alfaview software, the

amplified fragment sizes were estimated and discovered to be between 200 and 900 bp. Thus, the genetic relationship and degree of polymorphism of the various *T. cordifolia* samples were assessed using the illuminating OPA-01 primer as well as the sample obtained from Salem had the highest level of polymorphism.






Fig. 1. I and II demonstrate the samples underwent random amplification using the RAPD primers OPA01 and OPC2, which demonstrates the greatest polymorphism; III and IV demonstrate that samples underwent random amplification using the RAPD primers OPC14 and OPA03, which demonstrates the finest polymorphism; OPA-13, OPA-4, OPC-4, and OPC-1 RAPD primers, random amplification was performed in and the results indicated minimal polymorphism (V, VI, VII, and VIII)

*L (Ladder) Sample from Lane 1 comes from Thittacheri, Nagapattinam; Sample from lane 2 comes from Virudhachalam, Cuddalore; Sample from Lane 3 comes from Jayankondam, Ariyalur; Sample from Lane 4 comes from Kalavai, Ranipet; Sample from Lane 5 comes from Thalaivasal, Salem ; Sample from Lane 6 comes from Vedasandur, Dindigul; Sample from Lane 7 comes from Gudiyatham, Vellore; Sample from Lane 8 comes from Kinethukadavu, Coimbatore; Sample from Lane 9 comes from Vaniyampadi, Thirupathur; Sample from Lane 10 comes from Palani, Dindigul

Table 3. *T. cordifolia* sample RAPD analysis using primers

S. No.	CODES	No. of amplified alleles	No. of polimorphic alleles	No. of monomorphic alleles	polymorphic %	Amplification Status
1	OPA - 01	6	6	0	100	Amplified
2	OPC - 02	5	4	1	80	Amplified
3	OPC - 14	6	6	0	100	Amplified
4	OPA - 03	4	4	0	100	Amplified
5	OPA - 09	5	5	0	100	Amplified
6	OPC - 04	4	4	0	100	Amplified
7	OPC - 15	4	4	0	100	Amplified
8	OPA - 13	4	4	0	100	Amplified
9	OPA - 17	-	-	-	-	No Amplification
10	OPA - 20	-	-	-	-	No Amplification
11	OPG - 11	-	-	-	-	No Amplification
12	OPC - 11	-	-	-	-	No Amplification
13	OPG - 02	-	-	-	-	No Amplification
14	OPA - 04	-	-	-	-	No Amplification
15	TC - 01	-	-	-	-	No Amplification
16	TC - 02	-	-	-	-	No Amplification
17	TC - 03	-	-	-	-	No Amplification
18	OPG - 24	-	-	-	-	No Amplification

Using RAPD markers, the genetic diversity of 15 Bhubaneswar-collected *T. cordifolia* clones was evaluated. There were 40 decamer primers used. To determine the genetic links among 15 clones, 15 primers were utilised, and these primers

produced unique fragments of DNA [15]. Using RAPD markers, Shinde and Dhalwal [16] discovered limited amount of *T. cordifolia*'s genetic diversity. Twenty Western Himalayan *T. cordifolia* samples were chosen for genetic




Fig. 2. Ten samples' dendrogram demonstrating their genetic relatedness

analysis using RAPD markers. Out of the 120 decamer oligonucleotide primers employed, four RAPD primers only OPC-13, OPA-16, OPC-07 and OPC-05 exhibited minimal genetic variation of *T. cordifolia* [17]. 40 germplasm were gathered from different parts of India, including Delhi, Kerala, Himachal Pradesh, Assam, and Jammu & Kashmir. Three species of *Tinospora* were polymorphized by five out of nine primers [18]. The unweighted paired group method (UPGM) dendrogram produced by Ishnava and Mohan [19] demonstrated the extent of genetic diversity among several *T. cordifolia* germplasm.

Cluster analysis and Similarity matrix: Ten samples' dendrogram demonstrating their genetic relatedness shows on Fig. 2. The samples TCTV, TCNT, TCDP, and TCVG were discovered to be genetically related and to exhibit reduced polymorphism. However, sample TCCK was only distantly related to samples TCTV and TCNT and genetically distinct from samples TCDP and TCVG. Despite being distantly related to sample TCCV, TCDV and TCAJ shared genetic similarities. The samples TCTV and TCNT were closely linked to TCDV, TCAJ, and TCCV. There was no genetic overlap between TCCV and TCCK. With the most polymorphism, TCCV and TCCK were distantly linked to all other samples. When compared to other samples, TCST had the highest level of genetic diversity among the *T. cordifolia* accessions that had been gathered.

4. CONCLUSION

Ten accessions of *T. cordifolia* were gathered for the current study from diverse locations

throughout several districts of Tamil Nadu. Eight primers out of the 18 primers were capable of spotting differences between the 10 *T. cordifolia* samples that were obtained. Following analysis of RAPD, it was discovered that the DNA recovered from several *T. cordifolia* plants exhibited random amplification and various levels of genetic polymorphism. In the sample from Salem district the most genetic polymorphism was found.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Dubey NK, Kumar R, Tripathi P. Global promotion of herbal medicine: India's opportunity. *Current Science*. 2004; 86(1):37-41.
2. Foster BC, Arnason JT, Briggs CJ. Natural health products and drug disposition. *Annu. Rev. Pharmacol. Toxicol.* 2005; 45:203-26.
3. Singh A. A note on variation of active principles in Indian medicinal plants and TIM formulations. *Ethnobotanical leaflets*. 2008; (1):80.
4. Sharma A, Shanker C, Tyagi LK, Singh M, Rao CV. Herbal medicine for market potential in India: an overview. *Acad J Plant Sci*. 2008;1(2):26-36.
5. Singh SS, Pandey SC, Srivastava S, Gupta VS, Patro B. Chemistry, and medicinal properties of *Tinospora cordifolia*

- (Guduchi). Indian journal of pharmacology. 2003;35(2):83.
6. Kapoor LD. Handbook of Ayurvedic medicinal plants: Herbal reference library. CRC press; 2000.
 7. Bi D, Chen D, Khayatnezhad M, Hashjin ZS, Li Z, Ma Y. Molecular identification and genetic diversity in *Hypericum* L.: A high value medicinal plant using RAPD markers. Genetika. 2021;53(1):393-405.
 8. Khan S, Mirza KJ, Abdin MZ. Development of RAPD markers for authentication of medicinal plant *Cuscuta reflexa*. EurAsian Journal of Bio Sciences. 2010; 4.
 9. Mohapatra A, Rout GR. Identification and analysis of genetic variation among rose cultivars using random amplified polymorphic DNA. Zeitschrift für Naturforschung C. 2005;60(7-8):611-7.
 10. Britto DAJ, Raj TL, Petchimuthu K, Kumar PB, Sujin RM, Dharmar K. Inter population genetic variability in *Tinospora cordifolia* (Willd.) Miers ex Hook. F. & Thoms. (Menispermaceae), through RAPD marker. Acta Xaver. 2010;1:57-64.
 11. Aboul-Maaty NA, Oraby HA. Extraction of high-quality genomic DNA from different plant orders applying a modified CTAB-based method. Bulletin of the National Research Centre. 2019; 43(1):1-0.
 12. Srivastava P. *Tinospora cordifolia* (Amrita)- A miracle herb and lifeline to many diseases. International Journal of Medicinal and Aromatic Plants. 2011;1(2): 57-61.
 13. Rout GR. Direct plant regeneration from leaf explants of *Plumbago* species and its genetic fidelity through RAPD markers. Annals of applied biology. 2002;140(3): 305-13.
 14. Saha S, Dey ST, Adhikari S, Ghosh PD. In vitro multiple shoot regeneration and analysis of genetic fidelity of *Mentha piperita* L. Bionature. 2010;71-81.
 15. Rout GR. Identification of *Tinospora cordifolia* (Willd.) Miers ex Hook F. & Thomas Using RAPD Markers. Zeitschrift für Naturforschung C. 2006;61(1-2):118-22.
 16. Shinde VM, Dhalwal K. DNA fingerprinting of *Tinospora cordifolia* using RAPD analysis. J Global Pharm Technol. 2010; 2:38-42.
 17. Sharma R, Kumar V, Mohapatra T, Khandelwal V, Vyas GK. A simple and non-destructive method of direct-PCR for plant systems. Journal of Plant Biology. 2012; 55:114-22.
 18. Ahmad SM, Hoot SB, Qazi PH, Verma V. Phylogenetic patterns and genetic diversity of Indian *Tinospora* species based on chloroplast sequence data and cytochrome P450 polymorphisms. Plant systematics and evolution. 2009; 281:87-96.
 19. Ishnava K, Mohan JS. Assessment of genetic diversity in medicinal climber of *Tinospora cordifolia* (Willd.) Miers (Menispermaceae) from Gujarat, India. Asian Journal of Biotechnology. 2009;1(3): 93-103.

© 2023 Akhilraj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/99494>